

## PRODUCT MONOGRAPH

### **OLIMEL 3.3% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
3.3% w/v & 0.7% w/v, 11.5% w/v, 4% w/v

### **OLIMEL 4.4%**

Amino acids, dextrose, lipids  
Injectable Emulsion  
4.4% w/v, 14% w/v, 4% w/v

### **OLIMEL 4.4% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
4.4% w/v & 0.7% w/v, 14% w/v, 4% w/v

### **OLIMEL 5.7%**

Amino acids, dextrose, lipids  
Injectable Emulsion  
5.7% w/v, 11% w/v, 4% w/v

### **OLIMEL 5.7% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
5.7% w/v & 0.7% w/v, 11% w/v, 4% w/v

### **OLIMEL 7.6%**

Amino acids, dextrose, lipids  
Injectable Emulsion  
7.6% w/v, 7.3% w/v, 3.5% w/v

### **OLIMEL 7.6% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
7.6% w/v & 0.8% w/v, 7.3% w/v, 3.5% w/v

### **PeriOLIMEL 2.5% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
2.5% w/v & 0.4% w/v, 7.5% w/v, 3% w/v

## Intravenous Nutritive Supplements

Baxter Corporation  
Mississauga, Ontario L5N 0C2  
Canada

Date of Preparation:  
May 20, 2020

Submission Control No: 236356

Baxter, Oliclinomel, Olimel, and Periolimel are trademarks of Baxter International Inc.

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION.....</b>	<b>3</b>
<b>SUMMARY PRODUCT INFORMATION.....</b>	<b>4</b>
<b>INDICATIONS AND CLINICAL USE.....</b>	<b>5</b>
<b>CONTRAINDICATIONS.....</b>	<b>5</b>
<b>WARNINGS AND PRECAUTIONS.....</b>	<b>6</b>
<b>ADVERSE REACTIONS.....</b>	<b>11</b>
<b>DRUG INTERACTIONS.....</b>	<b>13</b>
<b>DOSAGE AND ADMINISTRATION.....</b>	<b>14</b>
<b>OVERDOSAGE.....</b>	<b>24</b>
<b>ACTION AND CLINICAL PHARMACOLOGY.....</b>	<b>24</b>
<b>STORAGE AND STABILITY.....</b>	<b>27</b>
<b>SPECIAL HANDLING INSTRUCTIONS.....</b>	<b>27</b>
<b>DOSAGE FORMS, COMPOSITION AND PACKAGING.....</b>	<b>32</b>
<b>PART II: SCIENTIFIC INFORMATION.....</b>	<b>37</b>
<b>PHARMACEUTICAL INFORMATION.....</b>	<b>37</b>
<b>CLINICAL TRIALS.....</b>	<b>41</b>
<b>DETAILED PHARMACOLOGY.....</b>	<b>42</b>
<b>TOXICOLOGY.....</b>	<b>42</b>
<b>REFERENCES.....</b>	<b>46</b>
<b>PART III: CONSUMER INFORMATION.....</b>	<b>50</b>

**OLIMEL 3.3% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
3.3% w/v & 0.7% w/v, 11.5% w/v, 4% w/v

**OLIMEL 4.4%**

Amino acids, dextrose, lipids  
Injectable Emulsion  
4.4% w/v, 14% w/v, 4% w/v

**OLIMEL 4.4% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
4.4% w/v & 0.7% w/v, 14% w/v, 4% w/v

**OLIMEL 5.7%**

Amino acids, dextrose, lipids  
Injectable Emulsion  
5.7% w/v, 11% w/v, 4% w/v

**OLIMEL 5.7% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
5.7% w/v & 0.7% w/v, 11% w/v, 4% w/v

**OLIMEL 7.6%**

Amino acids, dextrose, lipids  
Injectable Emulsion  
7.6% w/v, 7.3% w/v, 3.5% w/v

**OLIMEL 7.6% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
7.6% w/v & 0.8% w/v, 7.3% w/v, 3.5% w/v

**PeriOLIMEL 2.5% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
2.5% w/v & 0.4% w/v, 7.5% w/v, 3% w/v

PART I: HEALTH PROFESSIONAL INFORMATION

**SUMMARY PRODUCT INFORMATION**

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Intravenous	<p>Emulsion for Infusion /</p> <p>OLIMEL 3.3% E Amino acids <b>WITH</b> electrolytes, dextrose, lipids Injectable Emulsion 3.3% w/v &amp; 0.7% w/v, 11.5% w/v, 4% w/v</p> <p>OLIMEL 4.4% Amino acids, dextrose, lipids Injectable Emulsion 4.4% w/v, 14% w/v, 4% w/v</p> <p>OLIMEL 4.4% E Amino acids <b>WITH</b> electrolytes, dextrose, lipids Injectable Emulsion 4.4% w/v &amp; 0.7% w/v, 14% w/v, 4% w/v</p> <p>OLIMEL 5.7% Amino acids, dextrose, lipids Injectable Emulsion 5.7% w/v, 11% w/v, 4% w/v</p> <p>OLIMEL 5.7% E Amino acids <b>WITH</b> electrolytes, dextrose, lipids Injectable Emulsion 5.7% w/v &amp; 0.7% w/v, 11% w/v, 4% w/v</p> <p>OLIMEL 7.6% Amino acids, dextrose, lipids Injectable Emulsion 7.6% w/v, 7.3% w/v, 3.5% w/v</p> <p>OLIMEL 7.6% E Amino acids <b>WITH</b> electrolytes, dextrose, lipids Injectable Emulsion 7.6% w/v &amp; 0.8% w/v, 7.3% w/v, 3.5% w/v</p> <p>PeriOLIMEL 2.5% E Amino acids <b>WITH</b> electrolytes, dextrose, lipids Injectable Emulsion 2.5% w/v &amp; 0.4% w/v, 7.5% w/v, 3% w/v</p>	<p>Egg phosphatides</p> <p><i>For a complete listing see Dosage Forms, Composition and Packaging section.</i></p>

## **INDICATIONS AND CLINICAL USE**

OLIMEL/ PeriOLIMEL (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids) is indicated for parenteral nutrition for adults when oral or enteral nutrition is impossible, insufficient or contraindicated.

### **Geriatrics:**

There are no known differences in safety and effectiveness of parenteral nutrition formulations in the adult population based upon age.

### **Pediatrics:**

There have been no studies performed in the pediatric population.

## **CONTRAINDICATIONS**

The use of OLIMEL/ PeriOLIMEL (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids) is contraindicated in the following populations/situations:

- Known hypersensitivity to egg, soybean products, olive products or any of the active substances, excipients, or components of the container. For a complete listing, see the Dosage Forms, Composition and Packaging section of the product monograph.
- Known allergy to corn or corn products since the products contain corn-derived dextrose
- Patients with acute renal failure and without undergoing renal replacement therapy.
- Patients with severe liver failure or hepatic coma.
- Congenital abnormalities of amino acid metabolism
- Severe hyperlipidemia or severe disorders of lipid metabolism characterized by hypertriglyceridemia
- Hypertriglyceridemia-associated acute pancreatitis
- Severe hyperglycemia

Additional contraindications specific to OLIMEL / PeriOLIMEL formulations with electrolytes:

- Hyperkalemia (see **General** in **WARNINGS AND PRECAUTIONS**).

- Hypercalcaemia (see **General** in **WARNINGS AND PRECAUTIONS**).
- Hyperphosphatemia (see **General** in **WARNINGS AND PRECAUTIONS**)
- Hybernatremia.
- Hypermagnesemia.
- Ceftriaxone must not be administered simultaneously with intravenous calcium-containing solutions, including OLIMEL / PeriOLIMEL, through the same infusion line (e.g. via Y-site) because of the risk of precipitation of ceftriaxone-calcium salt. If the same infusion line is used for sequential administration, the line must be thoroughly flushed with a compatible fluid between infusions.

## **WARNINGS AND PRECAUTIONS**

### **General**

No substances or medicinal products other than the provided components of the products should be added to the bag without first confirming their compatibility and/or suitability to prevent formation of precipitates, destabilization of the lipid emulsion and/or overloading of electrolytes which may result in serious adverse reactions (see **DRUG INTERACTIONS** and **SPECIAL HANDLING INSTRUCTIONS**).

Due to presence of phosphate ion in OLIMEL / PeriOLIMEL products (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids), administration of these products may result in precipitation of calcium phosphate in patients with hyperphosphatemia, hypercalcaemia and /or co-administrated with a calcium ion-containing IV solution. Caution must be exercised when calcium-containing agents are added in the product since this may result in formation of precipitates which can lead to serious or even fatal reactions (See Respiratory subsection below and **SPECIAL HANDLING INSTRUCTIONS**).

The infusion must be stopped immediately if any signs or symptoms of an allergic reaction (such as fever, shivering, sweating, headache, skin rashes, or dyspnea) develop. Specific clinical monitoring is required when an intravenous nutrition infusion is started.

Infection and sepsis may occur as a result of improper use of intravenous catheters to administer parenteral formulations, poor maintenance of catheters or contaminated solutions. Immunosuppression and other conditions such as hyperglycemia, malnutrition and/or their underlying disease state may predispose patients to infectious complications. Careful symptomatic and laboratory monitoring for fever/chills, leukocytosis, technical complications with the access device, and hyperglycemia can help recognise early infections. The occurrence of septic complications can be

decreased with heightened emphasis on aseptic technique in catheter placement and maintenance as well as aseptic technique in nutritional formula preparation.

“Fat overload syndrome” has been reported with similar products. This may be caused by inappropriate administration (e.g. overdose and/or infusion rate higher than recommended, see **ADVERSE REACTIONS** and **OVERDOSAGE**); however, the signs and symptoms of this syndrome may also occur when the product is administered according to instructions. The reduced or limited ability to metabolize the lipids contained in OLIMEL / PeriOLIMEL may result in a fat overload syndrome. This syndrome is associated with a sudden deterioration in the patient's clinical condition and is characterized by hyperlipidemia, fever, jaundice, liver fatty infiltration (hepatomegaly), deteriorating liver function (hepatosplenomegaly), and hypoxia with or without respiratory insufficiency, anemia, leucopenia, thrombocytopenia, coagulation disorders and central nervous system manifestations (e.g. coma). These symptoms are usually reversible when the infusion of lipid emulsion is stopped.

Refeeding severely undernourished patients may result in the refeeding syndrome that is characterized by the shift of potassium, phosphorus, and magnesium intracellularly as the patients becomes anabolic. Thiamine deficiency and fluid retention may also develop. Careful monitoring and slowly increasing nutrient intakes while avoiding overfeeding can prevent these complications. This syndrome has been reported with similar products.

If the final mixture is hypertonic, it may cause irritation of the vein when administered into a peripheral vein.

While PeriOLIMEL may be administered through a peripheral or central vein, thrombophlebitis may develop if peripheral veins are used. The catheter insertion site must be monitored daily for local signs of thrombophlebitis.

OLIMEL must only be administered through a central vein.

Do not connect bags in series in order to avoid air embolism due to possible residual gas contained in the primary bag.

### **Carcinogenesis and Mutagenesis**

See **TOXICOLOGY** in Part II of the Product Monograph.

### **Extravasation**

Extravasation has been reported with the administration of OLIMEL / PeriOLIMEL.

## **Cardiovascular**

Use with caution in patients with pulmonary edema or heart failure. Fluid status should be closely monitored. The level of triglyceride should be monitored to avoid hypertriglyceridemia when administering OLIMEL / PeriOLIMEL in patients with acute myocardial infarction.

## **Endocrine and Metabolism**

Metabolic complications may occur if the nutrient intake is not adapted to the patient's requirements, or the metabolic capacity of any given dietary component is not accurately assessed. Adverse metabolic effects may arise from administration of inadequate or excessive nutrients or from inappropriate composition of an admixture for a particular patient's needs.

Serum triglycerides concentrations and the ability of the body to metabolize lipids must be monitored regularly. If a lipid metabolism abnormality is suspected, daily monitoring of serum triglycerides is recommended. Hypertriglyceridemia left untreated can lead to the development of pancreatitis, altered pulmonary function, and immune dysfunction.

Hypercholesterolemia may be caused by excessive amounts of phospholipids in the parenteral formula.

In the event of hyperglycemia, the infusion rate of OLIMEL / PeriOLIMEL must be adjusted and/or insulin administered.

### **Fructose**

This product may contain fructose as an impurity in the dextrose material. Exercise caution when this product is used in patients with hereditary fructose intolerance. In these patients, fructose may result in hypoglycemia, metabolic acidosis, liver toxicity which manifests as vomiting, nausea, sweating, jaundice, hemorrhage, seizures or coma or even death. The severity of the reactions is dependent on the amount and duration of fructose intake.

### **Hyperglycemia**

Rapid administration of dextrose solutions may produce substantial hyperglycemia which may result in or contribute to electrolyte losses, dehydration and hypovolemia due to osmotic diuresis and hyperosmolar syndrome. At certain clinical conditions it also may increase the risk of hyposmotic hyponatremia by shifting of intracellular water to extracellular space. Use with caution in critically ill patients in whom



hyperglycemia commonly occurs due to diabetes, impaired glucose intolerance, impaired fasting glucose, or is stress-induced. Hyperglycemia may increase the risk of cardiac complications, infection, systemic sepsis, acute renal failure and even death in certain clinical conditions, especially in acute stress conditions. In order to avoid hyperglycemia the infusion rate should not exceed the patient's ability to utilize glucose. To reduce the risk of hyperglycemia-associated complications, the infusion rate must be adjusted and/or insulin administered if blood glucose levels exceed levels considered acceptable for the individual patient.

### **Gastrointestinal**

Patients may develop nausea or diarrhea.

### **Hepatic/Biliary/Pancreatic**

Parenteral nutrition in general especially those containing amino acids, such as OLIMEL / PeriOLIMEL, should be used with caution in patients with preexisting liver disease or liver insufficiency. Liver function parameters should be closely monitored in these patients, and they should be monitored for possible symptoms of hyperammonemia.

Hepatobiliary disorders including cholestasis, hepatic steatosis, fibrosis and cirrhosis, possibly leading to hepatic failure, as well as cholecystitis and cholelithiasis are known to develop in some patients on parenteral nutrition. The etiology of these disorders is thought to be multifactorial and may differ between patients. Patients developing abnormal laboratory parameters or other signs of hepatobiliary disorders should be assessed early by a clinician knowledgeable in liver diseases in order to identify possible causative and contributory factors, and possible therapeutic and prophylactic interventions.

Elevated bilirubin and hepatic enzymes may occur in patients receiving parenteral nutrition and may result from excess administration of carbohydrate or lipid, lack of enteral nutrient stimulation, infection, or underlying disease.

### **Immune**

Hypersensitivity to the constituents of the parenteral nutrition formulation such as egg, soybean products, amino acids, olive products, or any of the active substances, excipients, or components of the containers may occur. See CONTRAINDICATIONS.

Since dextrose in OLIMEL / PeriOLIMEL products is derived from corn, these products should not be used in patients with known allergy to corn or corn products. See CONTRAINDICATIONS.

### **Renal**

Use with caution in patients with renal insufficiency. Fluid and electrolyte status should be closely monitored in these patients.

Severe water and electrolyte equilibration disorders, severe fluid overload states, and severe metabolic disorders should be corrected before starting the infusion.

### **Respiratory**

Pulmonary vascular precipitates causing pulmonary vascular emboli and pulmonary distress have been reported in patients receiving parenteral nutrition. In some cases, fatal outcomes have occurred. Excessive addition of calcium and phosphate increases the risk of the formation of calcium phosphate precipitates. Precipitates have been reported even in the absence of phosphate salt in the solution. Precipitation distal to the in-line filter and suspected precipitate formation in the blood stream have also been reported.

In addition to inspection of the solution, the infusion set and catheter should also periodically be checked for precipitates.

If signs of pulmonary distress occur, the infusion should be stopped and medical evaluation initiated.

Parenteral nutrition containing lipid emulsions should be given cautiously to patients with acute respiratory distress syndrome.

### **Special Populations**

#### **Pregnant Women:**

There are no adequate data on use of OLIMEL / PeriOLIMEL in pregnant women. Physicians should carefully consider the potential risks and benefits for each specific patient before prescribing OLIMEL / PeriOLIMEL.

#### **Nursing Women:**

There are no adequate data on use of OLIMEL / PeriOLIMEL in lactating women. Physicians should carefully consider the potential risks and benefits for each specific patient before prescribing OLIMEL / PeriOLIMEL.

### **Pediatrics:**

There have been no studies performed in the pediatric population.

### **Monitoring and Laboratory Tests**

Monitor water and electrolyte balance, serum osmolarity, serum triglycerides, acid/base balance, blood glucose, liver and kidney function, blood count, including platelets, and coagulation parameters throughout treatment. Daily monitoring is recommended during initiation of parenteral nutrition and until the patient and laboratory measurements are stable.

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

See **WARNINGS AND PRECAUTIONS**.

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

The safety and clinical efficacy of OLIMEL (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids) was assessed in a double-blind randomized controlled study over five days. Fifty-six (56) patients requiring parenteral nutrition were enrolled, of whom twenty-eight (28) were treated with OliClinomel (a triple-chamber parenteral nutrition product similar to OLIMEL, that contains the same olive oil/soybean oil lipid, a similar amino acid profile, and dextrose) and twenty-eight (28) were treated with OLIMEL. The goal of the study was to provide information on the safety and nutritional efficacy of OLIMEL in a clinical setting.

A total of fifty-three (53) adverse events occurred during treatment; twenty-nine (29) adverse events were observed in fourteen (14) patients in the OLIMEL group versus twenty-four (24) adverse events observed in eleven (11) patients in the OliClinomel

(control) group. Of the twenty-nine (29) adverse events observed in the OLIMEL group, seven (7) adverse events were designated as related to treatment. Of the twenty-four (24) adverse events observed in the OliClinomel (control) group, seven (7) patients presented with adverse events that were reported as related to treatment.

### Summary of Treatment-Related Adverse Drug Reactions in the OLIMEL Study

System Organ Class	Adverse Event	Reported Incidence by Treatment Group			
		OLIMEL (n=28) up to 40 mL/kg/day		OLICLINOMEL (n=28) up to 40 mL/kg/day	
		N*	%	N*	%
<b>Cardiac disorders</b>	Tachycardia	1	3.57	0	0.00
<b>Gastrointestinal disorders</b>	Abdominal pain	1	3.57	0	0.00
	Diarrhea	1	3.57	1	3.57
	Nausea	1	3.57	0	0.00
<b>Immune system disorders</b>	Hypersensitivity	0	0.00	1	3.57
<b>Investigations</b>	Blood alkaline phosphatase increased	0	0.00	1	3.57
	Gamma-glutamyltransferase increased	0	0.00	1	3.57
<b>Metabolism and nutrition disorders</b>	Decreased appetite	1	3.57	0	0.00
	Hypertriglyceridemia	1	3.57	0	0.00
<b>Renal and urinary disorders</b>	Azotemia	0	0.00	1	3.57
<b>Respiratory, thoracic and mediastinal disorders</b>	Respiratory failure	0	0.00	1	3.57
<b>Vascular disorders</b>	Hemodynamic instability	0	0.00	1	3.57
	Hypertension	1	3.57	0	0.00

\*Number of patients reporting the related event

### **Post-Market Adverse Drug Reactions**

In addition, the following adverse reactions have been reported in the postmarketing experience, listed by MedDRA System Organ Class (SOC), then by Preferred Term (PT) in order of severity.

GASTROINTESTINAL DISORDERS: Vomiting

SKIN AND SUBCUTANEOUS SKIN DISORDERS: Rash

GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS: Injection site extravasation, Pyrexia, Chills

### **Class Reactions**

The following adverse reactions have been reported with similar products:

Pruritus, Fat overload syndrome, Cholestasis, Elevated liver enzymes and Azotemia

Pulmonary vascular precipitates (pulmonary vascular emboli and pulmonary distress)  
(see WARNINGS AND PRECAUTIONS)

### **DRUG INTERACTIONS**

#### **Overview**

No interaction studies have been performed with OLIMEL / PeriOLIMEL (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids).

OLIMEL / PeriOLIMEL must not be administered simultaneously with blood through the same infusion tubing because of the risk of pseudoagglutination.

#### **Drug-Drug Interactions**

##### **Established or Potential Drug-Drug Interactions**

<b>&lt;Proper name&gt;</b>	<b>Ref</b>	<b>Effect</b>	<b>Clinical comment</b>
COUMADIN (or coumarin derivatives including warfarin)	T	Decreased anticoagulant effect	Soybean oil has a natural content of vitamin K that may counteract the anticoagulant activity of coumarin derivatives, including warfarin. Levels of vitamin K vary in the formulation. Caution is warranted and therapeutic monitoring of coagulation status is recommended.
Ceftriaxone	T	Precipitate forms	Ceftriaxone must not be administered with intravenous calcium-containing solutions, including OLIMEL / PeriOLIMEL, through the same infusion line (e.g. via Y-site) because of the risk of precipitation of ceftriaxone-calcium salt. If the same line is used for sequential administration, the line must be thoroughly flushed with a compatible fluid between infusions.

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

Due to the potassium content of OLIMEL / PeriOLIMEL special care should be taken in patients simultaneously treated with potassium sparing diuretics (e.g., amiloride, spironolactone, triamterene) with ACE inhibitors, angiotensin II receptor antagonists, or the immunosuppressants tacrolimus and cyclosporine in view of the risk of hyperkalemia. This only applies to products containing electrolytes.

Due to the risk of precipitation, OLIMEL / PeriOLIMEL should not be admixed with ampicillin or fosphenytoin.

Due to the presence of phosphate in OLIMEL / PeriOLIMEL and calcium ion in OLIMEL / PeriOLIMEL with electrolyte, adding calcium-containing and/or phosphate-containing products to the products may result in formation of calcium phosphate precipitates which may destabilize the emulsion status of the product (see SPECIAL HANDLING INSTRUCTIONS). Calcium phosphate precipitates and destabilized emulsion status of the products may result in serious health consequences.

### **Drug-Food Interactions**

No OLIMEL / PeriOLIMEL - food interaction studies have been performed.

### **Drug-Laboratory Interactions**

The lipids contained in this emulsion may interfere with the results of certain laboratory tests if the blood sample is taken before lipids are eliminated (these are generally eliminated after a period of 5 to 6 hours without receiving lipids). Potential assay interference associated with lipemia should be considered when interpreting the results of lipemic samples.

### **Drug-Lifestyle Interactions**

Interactions with lifestyle have not been evaluated.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

Dosing of OLIMEL / PeriOLIMEL (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids) is based upon protein, energy, and electrolyte requirements of the individual patient.

Some patients with renal and hepatic disease may benefit from reduced protein intakes.

Some patients with diabetes mellitus or glucose intolerance may benefit from reduced carbohydrate intake.

Patients with parenteral nutrition induced liver disease may benefit from reduction in lipid intake.

Patients with renal disease have decreased capacity to excrete potassium, magnesium, and phosphorus. Levels of these electrolytes should be carefully monitored in these patients.

### **Recommended Dose and Dosage Adjustment**

#### **Adults**

Due to its high osmolarity (1120-1360 mOsmol/L), OLIMEL must only be administered through a central vein.

Due to its low osmolarity (760 mOsmol/L), PeriOLIMEL can be administered through a peripheral or central vein.

The dosage depends on the patient's energy expenditure, clinical status, body weight, and ability to metabolize the constituents of OLIMEL / PeriOLIMEL, as well as additional energy or proteins/amino acids given orally/enterally; therefore, the bag size should be chosen accordingly.

The average daily requirements in adults are:

- Protein/amino acids: 0.16 to 0.35 g nitrogen /kg body weight (1 to 2 g of amino acids/kg), depending on the patient's nutritional status and degree of catabolic stress. Special populations may require up to 0.4 g nitrogen/kg body weight (2.5 g of amino acids/kg).
- Energy: 20 to 40 kcal/kg,
- Fluid: 20 to 40\* mL fluid /kg, or 1 to 1.5 mL per expended kcal.

\*This is dependent on not over feeding the patient calories or protein.

The maximum daily dose should not be exceeded. Due to the static composition of the multi-chamber bag, the ability to simultaneously meet all nutrient needs of the patient may not be possible. Clinical situations may exist where patients require amounts of nutrients varying from the composition of the static bag. In this situation the impact of

any volume (dose) adjustments must take into consideration the resultant effect this will have on the dosing of all other nutrient components of OLIMEL / PeriOLIMEL products. For example, patients may require greater than 0.2 mmol/kg/day of phosphate. In those situations, healthcare professionals may consider adjusting the volume (dose) of OLIMEL /PeriOLIMEL in order to meet these increased requirements.

The reconstituted emulsion provides the following for each of the formulations and bag sizes:

**PeriOLIMEL 2.5%E**

	<b>1000 mL</b>	<b>1500 mL</b>	<b>2000 mL</b>	<b>2500 mL</b>
Lipids	30 g	45 g	60 g	75 g
Amino acids	25.3 g	38.0 g	50.6 g	63.3 g
Nitrogen	4.0 g	6.0 g	8.0 g	10.0 g
Dextrose Anhydrous	75.0 g	112.5 g	150.0 g	187.5 g
Energy:				
Total calories approx.	700 kcal	1050 kcal	1400 kcal	1750 kcal
Non-amino acid calories approx.	600 kcal	900 kcal	1200 kcal	1500 kcal
Glucose calories	300 kcal	450 kcal	600 kcal	750 kcal
Lipid calories approx. <sup>(1)</sup>	300 kcal	450 kcal	600 kcal	750 kcal
Non-amino acid calories / nitrogen ratio	150 kcal/g	150 kcal/g	150 kcal/g	150 kcal/g
Glucose / lipid calories ratio	50/50	50/50	50/50	50/50
Lipid / total calories	43%	43%	43%	43%
Electrolytes:				
Sodium	21 mmol	31.5 mmol	42 mmol	52.5 mmol
Potassium	16.0 mmol	24.0 mmol	32.0 mmol	40.0 mmol
Magnesium	2.2 mmol	3.3 mmol	4.4 mmol	5.5 mmol
Calcium	2.0 mmol	3.0 mmol	4.0 mmol	5.0 mmol
Phosphate <sup>(2)</sup>	8.5 mmol	12.7 mmol	17.0 mmol	21.2 mmol
Acetate	27 mmol	41 mmol	55 mmol	69 mmol
Chloride	24 mmol	37 mmol	49 mmol	61 mmol



pH approx.	6.4	6.4	6.4	6.4
Osmolarity approx.	760 mosm/L	760 mosm/L	760 mosm/L	760 mosm/L

<sup>1</sup> Includes calories from purified egg phosphatides

<sup>2</sup> Includes phosphate provided by the lipid emulsion

**OLIMEL 3.3%E**

	<b>1500 mL</b>	<b>2000 mL</b>	<b>2500 mL</b>
Lipids	60 g	80 g	100 g
Amino acids	49.4g	65.8 g	82.3 g
Nitrogen	7.8 g	10.4 g	13.0 g
Dextrose Anhydrous	172.5 g	230.0 g	287.5 g
Energy:			
Total calories approx.	1490 kcal	1980 kcal	2480 kcal
Non-amino acid calories approx.	1290 kcal	1720 kcal	2150 kcal
Glucose calories	690 kcal	920 kcal	1150 kcal
Lipid calories approx. <sup>(1)</sup>	600 kcal	800 kcal	1000 kcal
Non-amino acid calories/nitrogen ratio	165 kcal/g	165 kcal/g	165 kcal/g
Glucose / lipid calories ratio	53/47	53/47	53/47
Lipid /total calories	47%	47%	47%
Electrolytes:			
Sodium	52.5 mmol	70.0 mmol	87.5 mmol
Potassium	45.0 mmol	60.0 mmol	75.0 mmol
Magnesium	6.0 mmol	8.0 mmol	10.0 mmol
Calcium	5.3 mmol	7.0 mmol	8.8 mmol
Phosphate <sup>(2)</sup>	22.5 mmol	30.0 mmol	37.5 mmol
Acetate	55 mmol	73 mmol	91 mmol
Chloride	68 mmol	90 mmol	113 mmol
pH approx.	6.4	6.4	6.4
Osmolarity approx.	1120 mosm/L	1120 mosm/L	1120 mosm/L

<sup>1</sup> Includes calories from purified egg phosphatides<sup>2</sup> Includes phosphate provided by the lipid emulsion

**OLIMEL 4.4%E and OLIMEL 4.4%**

	<b>1000 mL</b>	<b>1500 mL</b>	<b>2000 mL</b>
Lipids	40 g	60 g	80 g
Amino acids	44.3 g	66.4 g	88.6 g
Nitrogen	7.0 g	10.5 g	14.0 g
Dextrose Anhydrous	140.0 g	210.0 g	280.0 g
Energy:			
Total calories approx.	1140 kcal	1710 kcal	2270 kcal
Non-amino acid calories approx.	960 kcal	1440 kcal	1920 kcal
Glucose calories	560 kcal	840 kcal	1120 kcal
Lipid calories approx. <sup>(1)</sup>	400 kcal	600 kcal	800 kcal
Non-amino acid calories/ nitrogen ratio	137 kcal/g	137 kcal/g	137 kcal/g
Glucose / lipid calories ratio	58/42	58/42	58/42
Lipid / total calories	35%	35%	35%
Electrolytes <sup>(2)</sup> :			
Sodium	35.0 mmol	52.5 mmol	70.0 mmol
Potassium	30.0 mmol	45.0 mmol	60.0 mmol
Magnesium	4.0 mmol	6.0 mmol	8.0 mmol
Calcium	3.5 mmol	5.3 mmol	7.0 mmol
Phosphate <sup>(3)</sup>	15.0 (3.0) mmol	22.5 (4.5) mmol	30.0 (6.0) mmol
Acetate	45 (31) mmol	67 (46) mmol	89 (62) mmol
Chloride	45 mmol	68 mmol	90 mmol
pH approx.	6.4	6.4	6.4
Osmolarity approx.	1360 mosm/L (1220 mosm/L) <sup>(4)</sup>	1360 mosm/L (1220 mosm/L) <sup>(4)</sup>	1360 mosm/L (1220 mosm/L) <sup>(4)</sup>

<sup>1</sup> Includes calories from purified egg phosphatides

<sup>2</sup> Electrolytes are included in the OLIMEL 4.4% E presentation only. Values in parentheses for phosphate and acetate are the values that are present in the OLIMEL 4.4% (without electrolytes) formulation.

<sup>3</sup> Includes phosphate provided by the lipid emulsion, either formulation.

<sup>4</sup> Approximate osmolarity for OLIMEL 4.4% (without electrolytes).

**OLIMEL 5.7%E and OLIMEL 5.7%**

	<b>1000 mL</b>	<b>1500 mL</b>	<b>2000 mL</b>
Lipids	40 g	60 g	80 g
Amino acids	56.9 g	85.4 g	113.9 g
Nitrogen	9.0 g	13.5 g	18.0 g
Dextrose Anhydrous	110.0 g	165.0 g	220.0 g
Energy:			
Total calories approx.	1070 kcal	1600 kcal	2140 kcal
Non-amino acid calories approx.	840 kcal	1260 kcal	1680 kcal
Glucose calories	440 kcal	660 kcal	880 kcal
Lipid calories approx. <sup>(1)</sup>	400 kcal	600 kcal	800 kcal
Non-amino acid calories / nitrogen ratio	93 kcal/g	93 kcal/g	93 kcal/g
Glucose / lipid calories ratio	52/48	52/48	52/48
Lipid / total calories	37%	37%	37%
Electrolytes <sup>(2)</sup> :			
Sodium	35.0 mmol	52.5 mmol	70.0 mmol
Potassium	30.0 mmol	45.0 mmol	60.0 mmol
Magnesium	4.0 mmol	6.0 mmol	8.0 mmol
Calcium	3.5 mmol	5.3 mmol	7.0 mmol
Phosphate <sup>(3)</sup>	15.0 (3.0) mmol	22.5 (4.5) mmol	30.0 (6.0) mmol
Acetate	54 (40) mmol	80 (60) mmol	107 (80) mmol
Chloride	45 mmol	68 mmol	90 mmol
pH approx.	6.4	6.4	6.4
Osmolarity approx.	1310 mosm/L (1170 mosm/L) <sup>(4)</sup>	1310 mosm/L (1170 mosm/L) <sup>(4)</sup>	1310 mosm/L (1170 mosm/L) <sup>(4)</sup>

<sup>1</sup> Includes calories from purified egg phosphatides

<sup>2</sup> Electrolytes are included in the OLIMEL 5.7% E presentation only. Values in parentheses for phosphate and acetate are the values that are present in the OLIMEL 5.7% (without electrolytes) formulation.

<sup>3</sup> Includes phosphate provided by the lipid emulsion, either formulation.

<sup>4</sup> Approximate osmolarity for OLIMEL 5.7% (without electrolytes).

**OLIMEL 7.6%E and OLIMEL 7.6%**

	<b>650 mL</b>	<b>1000 mL</b>	<b>1500 mL</b>	<b>2000 mL</b>
Lipids	22.8 g	35.0 g	52.5 g	70.0 g
Amino acids	49.4 g	75.9 g	113.9 g	151.9 g
Nitrogen	7.8 g	12.0 g	18.0 g	24.0 g
Dextrose Anhydrous	47.7 g	73.3 g	110.0 g	146.7 g
Energy:				
Total calories approx.	620 kcal	950 kcal	1420 kcal	1900 kcal
Non-amino acid calories approx.	420 kcal	640 kcal	960 kcal	1280 kcal
Glucose calories	190 kcal	290 kcal	430 kcal	580 kcal
Lipid calories approx. <sup>(1)</sup>	230 kcal	350 kcal	530 kcal	700 kcal
Non-amino acid calories / nitrogen ratio	53 kcal/g	53 kcal/g	53 kcal/g	53 kcal/g
Glucose / lipid calories ratio	45/55	45/55	45/55	45/55
Lipid / total calories	37%	37%	37%	37%
Electrolytes <sup>(2)</sup> :				
Sodium	22.8 mmol	35.0 mmol	52.5 mmol	70.0 mmol
Potassium	19.5 mmol	30.0 mmol	45.0 mmol	60.0 mmol
Magnesium	2.6 mmol	4.0 mmol	6.0 mmol	8.0 mmol
Calcium	2.3 mmol	3.5 mmol	5.3 mmol	7.0 mmol
Phosphate <sup>(3)</sup>	9.5 (1.7) mmol	15.0 (2.6) mmol	21.9 (3.9) mmol	29.2 (5.2) mmol
Acetate	46 (35) mmol	70 (54) mmol	105 (80) mmol	140 (107) mmol
Chloride	30 mmol	45 mmol	68 mmol	90 mmol
pH approx.	6.4	6.4	6.4	6.4
Osmolarity approx.	1270 mosm/ L (1130 mosm/L) <sup>(4)</sup>	1270 mosm/L (1130 mosm/L) <sup>(4)</sup>	1270 mosm/L (1130 mosm/L) <sup>(4)</sup>	1270 mosm/L (1130 mosm/L) <sup>(4)</sup>

<sup>1</sup> Includes calories from purified egg phosphatides

<sup>2</sup> Electrolytes are included in the OLIMEL 7.6% E presentation only. Values in parentheses for phosphate and acetate are the values that are present in the OLIMEL 7.6% (without electrolytes) formulation.

<sup>3</sup> Includes phosphate provided by the lipid emulsion, either formulation.

<sup>4</sup> Approximate osmolarity for OLIMEL 7.6% (without electrolytes).

As a general rule, daily doses of 2 g/kg of amino acids and/or 7 g/kg of dextrose and/or 2 g/kg of lipids and/or 40 kcal/kg and/or 40 mL fluid/kg should not be exceeded, except in particular cases when specific demands exceed the usual recommendations. The first of these doses to be reached sets the maximum daily dose.

The OLIMEL / PeriOLIMEL formulations may be limited by fluid (40 ml/kg/day), energy (40 kcal/kg/day total calories) or amino acids intakes (2g/kg/day). In Continuous Renal Replacement (CRRT) and patients with morbid obesity, the maximal daily dose is defined by amino acids intake. The maximal daily dose delivers the following:

	<b>PeriOLIMEL 2.5%E</b>	<b>OLIMEL 3.3%E</b>	<b>OLIMEL 4.4% E and OLIMEL 4.4%</b>	<b>OLIMEL 5.7% E and OLIMEL 5.7%</b>	<b>OLIMEL 7.6% E and OLIMEL 7.6%</b>	<b>OLIMEL 7.6% E and OLIMEL 7.6%**</b>
Fluid volume (mL/kg/day)	40	40	35	35	26	33
Energy (kcal/kg/day)	28	40	40	37	24.7	31.2
Amino Acids (g/kg/day)	1	1.3	1.5	2	2.0	2.5***
Dextrose (g/kg/day)	3	4.6	4.9	3.9	1.9	2.4
Lipid (g/kg/day)	1.2	1.6	1.4	1.4	0.9	1.2
Sodium (mmol/kg/day)	0.8	1.4	1.2*	1.2*	0.9*	1.2*
Potassium (mmol/kg/day)	0.6	1.2	1.1*	1.1*	0.8*	1.0*

\* Only for electrolyte (E) formulations

\*\* In Continuous Renal Replacement (CRRT) and patients with morbid obesity

\*\*\* In patients with morbid obesity, dosing is based on ideal body weight (IBW)

The maximal infusion rates for the average patient receiving OLIMEL / PeriOLIMEL formulations over a 24 hour administration period are as follows:

<b>Formulation</b>	<b>Maximal infusion rate</b>	<b>Amino acids (g/kg/hour)</b>	<b>Dextrose (g/kg/hour)</b>	<b>Lipids (g/kg/hour)</b>
PeriOLIMEL 2.5% E	3.2 mL/kg/hour	0.08	0.24	0.1
OLIMEL 3.3%E	2.1 mL/kg/hour	0.07	0.24	0.08
OLIMEL 4.4% and OLIMEL 4.4% E	1.7 mL/kg/hour	0.08	0.24	0.07
OLIMEL 5.7% and OLIMEL 5.7% E	1.8 mL/kg/hour	0.1	0.19	0.07
OLIMEL 7.6% and OLIMEL 7.6% E	1.3 mL/kg/hour	0.10	0.10	0.05

The flow rate should be increased gradually during the first hour. The administration flow rate must be adjusted to take into account the dose being administered, the daily volume intake and the duration of the infusion (see **OVERDOSAGE**).

### **Missed Dose**

In the event of a missed dose, the infusion should be restarted at the recommended dose and flow rate. Doses should NOT be doubled.

### **Administration**

For instructions for preparation and handling of the emulsion for infusion see SPECIAL HANDLING INSTRUCTIONS.

PeriOLIMEL 2.5% E can be administered through a peripheral or central vein due to its lower osmolarity. OLIMEL must be delivered through a central vein. (See WARNINGS AND PRECAUTIONS.)

The recommended duration of infusion for a parenteral nutrition bag is between 12 and 24 hours. Treatment with parenteral nutrition may be continued for as long as is required by the patient's clinical conditions.

Monitoring of laboratory and clinical parameters is recommended (see WARNINGS AND PRECAUTIONS and Monitoring and Laboratory Tests).

The compatibility with solutions administered simultaneously via a common end section must be ensured.

### **Additions:**

Although there is a natural content of trace elements and vitamins in the product, the levels of such substances may not be sufficient to meet body's requirements and need to be supplemented in sufficient quantities to meet individual patient requirements and to prevent deficiencies from developing. However, no addition of any other substances or medicinal products to the bag should be made without first confirming their compatibility and/or suitability to prevent formation of precipitates, destabilization of the lipid emulsion and/or overloading of electrolytes which may result in serious adverse reactions (see DRUG INTERACTIONS and SPECIAL HANDLING INSTRUCTIONS).

For OLIMEL / PeriOLIMEL, calcium and phosphate ratios must be considered. Excess addition of calcium ion and/or inorganic and/or organic phosphate may result in the formation of calcium phosphate precipitates.

Iron ions should NOT be added to the bag since they may destabilize the lipid emulsion.

The electrolyte-containing OLIMEL / PeriOLIMEL products contain calcium ions which pose additional risk of coagulation precipitated in citrate anticoagulated/preserved blood or components.

## **OVERDOSAGE**

For suspected cases of drug overdose, contact the regional poison control centre.

In the event of inappropriate administration (overdose and/or infusion rate higher than recommended), nausea, vomiting, chills, headache, hot flush, hyperhidrosis and electrolyte disturbances and signs of hypervolemia or acidosis may occur and result in severe or fatal consequences. In such situations the infusion must be stopped immediately. If medically appropriate, further intervention may be indicated.

Hyperglycemia, glucosuria, and heperosmolar syndrome may develop if dextrose infusion rate exceeds clearance.

In some serious cases, hemodialysis, hemofiltration or hemo-diafiltration may be necessary.

The reduced or limited ability to metabolize lipids may result in fat overload syndrome, the results of which are usually reversible after infusion of the lipid emulsion is stopped. (See **WARNINGS AND PRECAUTIONS** section)

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Amino acids administered intravenously are directly bioavailable to the body. The amino acids are transported into tissues by active transporters where they supply substrate for protein or peptide synthesis, act as regulators of various enzymes and genes, or are converted to other bioactive compounds (such as nitric oxide, glutathione, and gamma amino butyric acid). Thus, amino acids have both structural and regulatory roles within the body. Amino acids are metabolized in the liver and small quantities may be excreted unchanged through the kidneys. Nitrogen waste products from amino acid metabolism are converted to urea and excreted via the kidneys.

Dextrose, as glucose, is taken up and metabolized by all cells in the body and represents the primary energy source for the body. Glucose can be stored in the body in the form of glycogen (primarily in liver and skeletal muscle). Glucose may also be converted to fatty acids and stored as triglycerides in fat tissue. Circulating glucose levels are



regulated by the interplay between insulin and glucagon, with lesser contributions from catecholamines, growth hormone, and glucocorticoids.

Fatty acids (lipid) are important energy sources for the body. The human body cannot synthesize omega-6 (linoleic acid and derivatives) or omega-3 ( $\alpha$ -linolenic acid and derivatives) polyunsaturated fatty acids and requires these from the diet. Fatty acids are also important as substrates for membranes, precursors for bioactive molecules (such as prostaglandins), and as regulators of gene expression.

Electrolytes have many important functions in the body and are required by all cells in the body. Sodium is the primary regulator of extracellular volume and the major contributor to extracellular osmolality. Potassium is the primary determinant of the membrane electrical potential and important for protein synthesis. Calcium is essential for bone structure and function and is an important regular of cellular signalling and enzyme activity. Phosphorus is important for structure and function of many compounds, including the energy compounds, adenosine triphosphate and creatine phosphate. Magnesium is a regulator of many enzymes and a structural component of bone.

### **Pharmacodynamics**

The content of nitrogen (amino acids) in OLIMEL / PeriOLIMEL (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids) and energy (dextrose and triglycerides) enables maintenance of an adequate nitrogen/energy balance. Nitrogen and energy are required for normal functioning of all cells in the body, and are important for protein synthesis, growth, wound healing, immune function, muscle function, and many other cellular activities.

The amino acid solution contains 17 amino acids (including 8 essential amino acids), which are indispensable for protein synthesis. All the amino acids used in OLIMEL / PeriOLIMEL are in the optical L-form, except glycine, whose molecule has no chiral center and is not optically active. The optical L-form is known to be more compatible to human biochemistry. Amino acids also represent an energy source, their oxidation resulting in excretion of nitrogen in the form of urea.

The amino acids profile is as follows:

- Essential amino acids/total amino acids: 44.8%
- Branched-chain amino acids/total amino acids: 18.3%.

Electrolytes are required for normal function of all cells in the body. These include cardiovascular, neurologic, respiratory, renal, and hepatic functions.

The lipid emulsion included in OLIMEL / PeriOLIMEL, is a mixture of refined olive oil and refined soybean oil (ratio 80/20), with the following approximate distribution of fatty acids:

- 15% saturated fatty acids (SFA)
- 65% monounsaturated fatty acids (MUFA)
- 20% polyunsaturated essential fatty acids (PUFA)

The phospholipid/triglyceride ratio is 0.06. The moderate essential fatty acid (EFA) content may improve utilization of infused essential fatty acids for synthesis of higher derivative fatty acids.

Olive oil contains significant amounts of alpha-tocopherol that contributes to vitamin E status.

The carbohydrate source is dextrose. Dextrose (glucose) is the primary source of energy in the body.

### **Pharmacokinetics**

Absorption: Not applicable as this drug is given intravenously.

Distribution: The constituents of the formulation are distributed to all cells in the body.

The ingredients of OLIMEL / PeriOLIMEL (amino acids, electrolytes, dextrose, lipids) are distributed, metabolized and eliminated in the same manner as if they had been administered individually.

The pharmacokinetic properties of the amino acids administered intravenously are essentially the same as those of amino acids supplied by oral feeding. Amino acids from food proteins, however, first pass through the portal vein before reaching the systemic circulation.

The elimination rate of lipid emulsions depends on particle size, fatty acid composition, apolipoprotein content of the lipid globules, lipoprotein lipase activity, and hepatic lipase activity. The maximal removal capacity (K1) for the lipid emulsion found for OLIMEL / PeriOLIMEL in normal volunteers is  $176 \pm 16$  mg/kg/hr. In the emulsion

contained in OLIMEL / PeriOLIMEL, the size of the lipid particles is close to that of chylomicrons and this emulsion therefore has a similar elimination rate.

**Metabolism and excretion:** Amino acids, dextrose, and triglycerides are metabolized by all cells in the body. Nitrogen waste is converted to urea in the liver and excreted by the kidneys. Glucose and triglycerides are metabolized to carbon dioxide and excreted by the lungs. Electrolytes are not metabolized. They are stored in the body or excreted by the liver, intestines, kidneys, or skin.

### **Special Populations and Conditions**

Pharmacokinetic data have not been obtained in special patient populations or conditions.

### **STORAGE AND STABILITY**

Do not freeze. Store the unmixed product in the overpouch at 15°C to 30°C.

#### Shelf life after reconstitution:

It is recommended that the product be used immediately after the non-permanent seals between the 3 chambers have been opened.

#### Shelf life after addition of supplements (electrolytes, trace elements, vitamins; see **SPECIAL HANDLING INSTRUCTIONS**):

Addition of supplements must take place under controlled and validated aseptic conditions. From a microbiological point of view, any admixture should be used immediately. If not used immediately, storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C to 8°C.

### **SPECIAL HANDLING INSTRUCTIONS**

#### To open

Remove the protective overpouch.

Discard the oxygen absorber.

Confirm the integrity of the bag and of the non-permanent seals. Use only if the bag is not damaged, if the non-permanent seals are intact (i.e. no mixture of the contents of the three chambers), if the amino acids solution and the dextrose solution are clear,

colourless or slightly yellow, practically free of visible particles, and if the lipid emulsion is a homogeneous liquid with a milky appearance.

#### Mixing the solutions and the emulsion

Ensure that the product is at room temperature when breaking the non-permanent seals.

Manually roll the bag onto itself, starting at the top of the bag (hanger end). The non-permanent seals will disappear from the side near the inlets. Continue to roll until the seals are open along approximately half of their length.

Mix by inverting the bag at least 3 times.

The appearance after reconstitution is a homogeneous milk-like emulsion.

#### Additions

Additions must be performed under aseptic conditions and by qualified personnel.

The capacity of the bag is sufficient to enable additions of nutritional and medicinal substances such as vitamins, electrolytes and trace elements. However, other than the provided components of the product, no substance should be added to the bag without first confirming its compatibility and suitability as stated below in order to prevent formation of precipitates, destabilization of the lipid emulsion and/or overloading of electrolytes which may result in serious adverse reactions.

Any addition (including vitamins) may be made into the reconstituted mixture (after the non-permanent seals have been opened and after the contents of the three chambers have been mixed). Iron ions should NOT be added to the bag since they may destabilize the lipid emulsion of the product.

Vitamins may also be added into the dextrose chamber before the mixture is reconstituted (before opening the non-permanent seals and before mixing the solutions and the emulsion).

When making additions to PeriOLIMEL, the final osmolarity of the mixture should be measured before administration via a peripheral vein.

OLIMEL/PERIOLIMEL formulations may be supplemented with electrolytes. Electrolyte supplementation should be dictated by the patient's clinical needs and should not exceed nutritional guidelines. When making additions to the formulations containing electrolytes, the amount of electrolytes already present in the bag should be

taken into account. When calcium-containing and/or phosphate-containing products are added to the products, caution must be exercised to prevent formation of calcium phosphate precipitates in the reconstituted mixture AND/OR in the body after administration. The maximal total levels for sodium, magnesium, potassium and calcium listed in the tables below were demonstrated by stability data, but **should not be considered as dosage recommendations** since the safety of the resulting mixture has not been adequately studied in clinical settings. If the level of calcium and/or phosphate in the reconstituted mixture exceeds the level shown in the following tables, the stability of the mixture may be compromised due to formation of calcium phosphate precipitates.

**Additions to PeriOLIMEL 2.5% E:**

Per 1000 mL			
	Included level	Maximal further addition <sup>(4)</sup>	Maximal total level <sup>(4)</sup>
Sodium	21 mmol	129 mmol	150 mmol
Potassium	16 mmol	134 mmol	150 mmol
Magnesium	2.2 mmol	3.4 mmol	5.6 mmol
Calcium	2.0 mmol	3.0 (1.5 <sup>(2)</sup> ) mmol	5.0 (3.5 <sup>(2)</sup> ) mmol
Inorganic Phosphate	0 mmol	8.0 mmol	8.0 mmol
Organic Phosphate	8.5 mmol <sup>(1)</sup>	- <sup>(3)</sup>	- <sup>(3)</sup>

(1) Including phosphate provided by the lipid emulsion

(2) Value in condition where inorganic phosphate is added at the maximal addition level.

(3) Organic Phosphate as a single entity may not be currently available in Canada.

(4) The values are based on stability data and should not be considered as dose recommendations.

**Additions to OLIMEL 3.3% E, OLIMEL 4.4% E, OLIMEL 5.7% E:**

Per 1000 mL			
	Included level	Maximal further addition <sup>(4)</sup>	Maximal total level <sup>(4)</sup>
Sodium	35 mmol	115 mmol	150 mmol
Potassium	30 mmol	120 mmol	150 mmol
Magnesium	4.0 mmol	1.6 mmol	5.6 mmol
Calcium	3.5 mmol	1.5 (0.0 <sup>(2)</sup> ) mmol	5.0 (3.5 <sup>(2)</sup> )mmol
Inorganic Phosphate	0 mmol	3.0 mmol	3.0 mmol
Organic Phosphate	15 mmol <sup>(1)</sup>	- <sup>(3)</sup>	- <sup>(3)</sup>

- <sup>(1)</sup> Including phosphate provided by the lipid emulsion  
<sup>(2)</sup> Value in condition where inorganic phosphate is added at the maximal addition level.  
<sup>(3)</sup> Organic Phosphate as a single entity may not be currently available in Canada.  
<sup>(4)</sup> The values are based on stability data and should not be considered as dose recommendations.

**Additions to OLIMEL 4.4% and OLIMEL 5.7%:**

Per 1000 mL			
	Included level	Maximal further addition <sup>(4)</sup>	Maximal total level <sup>(4)</sup>
Sodium	0 mmol	150 mmol	150 mmol
Potassium	0 mmol	150 mmol	150 mmol
Magnesium	0 mmol	5.6 mmol	5.6 mmol
Calcium	0 mmol	5.0 (3.5 <sup>(2)</sup> ) mmol	5.0 (3.5 <sup>(2)</sup> ) mmol
Inorganic Phosphate	0 mmol	8.0 mmol	8.0 mmol
Organic Phosphate	3 mmol <sup>(1)</sup>	- <sup>(3)</sup>	- <sup>(3)</sup>

- <sup>(1)</sup> Including phosphate provided by the lipid emulsion  
<sup>(2)</sup> Value in condition where inorganic phosphate is added at the maximal addition level.  
<sup>(3)</sup> Organic Phosphate as a single entity may not be currently available in Canada.  
<sup>(4)</sup> The values are based on stability data and should not be considered as dose recommendations.

**Additions to OLIMEL 7.6 %:**

Per 1000 mL			
	Included level	Maximal further addition <sup>(4)</sup>	Maximal total level <sup>(4)</sup>
Sodium	0 mmol	150 mmol	150 mmol
Potassium	0 mmol	150 mmol	150 mmol
Magnesium	0 mmol	5.6 mmol	5.6 mmol
Calcium	0 mmol	5.0 (3.5 <sup>(2)</sup> ) mmol	5.0 (3.5 <sup>(2)</sup> ) mmol

Inorganic Phosphate	0 mmol	10.0 mmol	10.0 mmol
Organic Phosphate	3 mmol <sup>(1)</sup>	22 mmol <sup>(3)</sup>	25 mmol <sup>(3)</sup>

<sup>(1)</sup> Including phosphate provided by the lipid emulsion

<sup>(2)</sup> Value in condition where inorganic phosphate is added at the maximal addition level.

<sup>(3)</sup> Organic Phosphate as a single entity may not be currently available in Canada.

<sup>(4)</sup> The values are based on stability data and should not be considered as dose recommendations

### **Additions to OLIMEL 7.6 % E:**

<b>Per 1000 mL</b>			
	<b>Included level</b>	<b>Maximal further addition<sup>(4)</sup></b>	<b>Maximal total level<sup>(4)</sup></b>
Sodium	35 mmol	115 mmol	150 mmol
Potassium	30 mmol	120 mmol	150 mmol
Magnesium	4 mmol	1.6 mmol	5.6 mmol
Calcium	3.5 mmol	1.5 (0.0 <sup>(2)</sup> ) mmol	5.0 (3.5 <sup>(2)</sup> ) mmol
Inorganic Phosphate	0 mmol	10.0 mmol	10.0 mmol
Organic Phosphate	15 mmol <sup>(1)</sup>	10 mmol <sup>(3)</sup>	25 mmol <sup>(3)</sup>

<sup>(1)</sup> Including phosphate provided by the lipid emulsion

<sup>(2)</sup> Value in condition where inorganic phosphate is added at the maximal addition level.

<sup>(3)</sup> Organic Phosphate as a single entity may not be currently available in Canada.

<sup>(4)</sup> The values are based on stability data and should not be considered as dose recommendations

Trace elements and vitamins:

Stability has been demonstrated up to the recommended daily dose.

Trace elements and vitamins: Stability has been demonstrated with commercially available preparations of vitamins and trace elements (containing up to 1 mg of iron).

Compatibility for other additives is available upon request.

To perform an addition:

- Aseptic conditions must be observed.
- Prepare the injection site of the bag.
- Puncture the injection site and inject the additives using an injection needle or a reconstitution device.
- Mix content of the bag and the additives.

Preparation of the infusion

Only administration sets and administration lines made from DEHP-free should be used.

Aseptic conditions must be observed.

Suspend the bag.

Remove the plastic protector from the administration outlet.

Firmly insert the spike of the infusion set into the administration outlet.

### Administration

For single use only. Aseptic conditions must be observed. When additions are made with inorganic phosphate, 1.2 micron filters should be used.

Only administer the product after the non-permanent seals between the three chambers have been broken and the contents of the three chambers have been mixed.

Ensure that the final emulsion for infusion does not show any evidence of phase separation.

After opening the bag, the contents should be used immediately and should not be stored for a subsequent infusion. Do not reconnect any partially used bag.

Do not connect in series in order to avoid the possibility of air embolism due to gas contained in the first bag.

Any unused product or waste material and all necessary devices must be discarded.

### **DOSAGE FORMS, COMPOSITION AND PACKAGING**

OLIMEL / PeriOLIMEL (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids) is presented in the form of a 3-chamber bag. Each bag contains a sterile dextrose (glucose) solution with or without calcium, a sterile amino acid solution with or without other electrolytes and a sterile lipid emulsion as described below:

<b>PeriOLIMEL 2.5% E</b>	<b>Content per bag</b>			
	<b>1000 mL</b>	<b>1500 mL</b>	<b>2000 mL</b>	<b>2500 mL</b>
18.75% Dextrose solution (18.75 g/100 mL)	400 mL	600 mL	800 mL	1000 mL
6.3% Amino acid solution (6.3 g/100 mL)	400 mL	600 mL	800 mL	1000 mL
15% Lipid emulsion (15 g/ 100 mL)	200 mL	300 mL	400 mL	500 mL



**OLIMEL 3.3% E****Content per bag**

	<b>1500 mL</b>	<b>2000 mL</b>	<b>2500 mL</b>
28.75% Dextrose solution (28.75 g/100 mL)	600 mL	800 mL	1000 mL
8.2% Amino acid solution (8.2 g/100 mL)	600 mL	800 mL	1000 mL
20% Lipid emulsion (20 g/100 mL)	300 mL	400 mL	500 mL

**OLIMEL 4.4% E and OLIMEL 4.4%****Content per bag**

	<b>1000 mL</b>	<b>1500 mL</b>	<b>2000 mL</b>
35% Dextrose solution (35 g/100 mL)	400 mL	600 mL	800 mL
11.1% Amino acid solution (11.1 g/100 mL)	400 mL	600 mL	800 mL
20% Lipid emulsion (20 g/100 mL)	200 mL	300 mL	400 mL

**OLIMEL 5.7% E and OLIMEL 5.7%****Content per bag**

	<b>1000 mL</b>	<b>1500 mL</b>	<b>2000 mL</b>
27.5% Dextrose solution (27.5 g/100 mL)	400 mL	600 mL	800 mL
14.2% Amino acid solution (14.2 g/100 mL)	400 mL	600 mL	800 mL
20% Lipid emulsion (20 g/100 mL)	200 mL	300 mL	400 mL

**OLIMEL 7.6% E and OLIMEL 7.6%****Content per bag**

	<b>650 mL</b>	<b>1000 mL</b>	<b>1500 mL</b>	<b>2000 mL</b>
27.5% Dextrose solution (27.5 g/100 mL)	173 mL	267 mL	400 mL	533 mL
14.2% Amino acid solution (14.2 g/100 mL)	347 mL	533 mL	800 mL	1067 mL
17.5% Lipid emulsion (17.5 g/100 mL)	130 mL	200 mL	300 mL	400 mL

After mixing the contents of the three chambers, the composition per liter of the reconstituted emulsion is given in the following table for each formulation:

Active substances	Composition per Liter of Reconstituted Emulsion							
	PeriOLIMEL 2.5% E	OLIMEL 3.3% E	OLIMEL 4.4% E	OLIMEL 4.4%	OLIMEL 5.7% E	OLIMEL 5.7%	OLIMEL 7.6 % E	OLIMEL 7.6 %
Refined olive oil + refined soybean oil <sup>1</sup>	30.00 g	40.00 g	40.00 g	40.00 g	40.00 g	40.00 g	35.00 g	35.00 g
L-Alanine	3.66 g	4.76 g	6.41 g	6.41 g	8.24 g	8.24 g	10.99 g	10.99 g
L-Arginine	2.48 g	3.22 g	4.34 g	4.34 g	5.58 g	5.58 g	7.44 g	7.44 g
L-Aspartic acid	0.73 g	0.95 g	1.28 g	1.28 g	1.65 g	1.65 g	2.20 g	2.20 g
L-Glutamic acid	1.26 g	1.64g	2.21 g	2.21 g	2.84 g	2.84 g	3.79 g	3.79 g
Glycine	1.76 g	2.28 g	3.07 g	3.07 g	3.95 g	3.95 g	5.26 g	5.26 g
L-Histidine	1.51 g	1.96 g	2.64 g	2.64 g	3.40 g	3.40 g	4.53 g	4.53 g
L-Isoleucine	1.26 g	1.64 g	2.21 g	2.21 g	2.84 g	2.84 g	3.79 g	3.79 g
L-Leucine	1.76 g	2.28 g	3.07 g	3.07 g	3.95 g	3.95 g	5.26 g	5.26 g
L-Lysine acetate (equivalent to Lysine)	2.81 g (1.99 g)	3.65 g (2.59 g)	4.88 g (3.48 g)	4.88 g (3.48 g)	6.32 g (4.48 g)	6.32 g (4.48 g)	8.43 g (5.97 g)	8.43 g (5.97 g)
L-Methionine	1.26 g	1.64 g	2.21 g	2.21 g	2.84 g	2.84 g	3.79 g	3.79 g
L-Phenylalanine	1.76 g	2.28 g	3.07 g	3.07 g	3.95 g	3.95 g	5.26 g	5.26 g
L-Proline	1.51 g	1.96g	2.64 g	2.64 g	3.40 g	3.40 g	4.53 g	4.53 g
L-Serine	1.00 g	1.30 g	1.75 g	1.75 g	2.25 g	2.25 g	3.00 g	3.00 g
L-Threonine	1.26 g	1.64 g	2.21 g	2.21 g	2.84 g	2.84 g	3.79 g	3.79 g
L-Tryptophan	0.42 g	0.55 g	0.74 g	0.74 g	0.95 g	0.95 g	1.26 g	1.26 g
L-Tyrosine	0.06 g	0.08 g	0.11 g	0.11 g	0.15 g	0.15 g	0.20 g	0.20 g
L-Valine	1.62 g	2.10 g	2.83 g	2.83 g	3.64 g	3.64 g	4.86 g	4.86 g
Sodium acetate, trihydrate	1.16 g	1.50 g	1.50 g	--	1.50 g	--	1.50 g	--
Sodium glycerophosphate, hydrated	1.91 g	3.67 g	3.67 g	--	3.67 g	--	3.67 g	--
Potassium chloride	1.19 g	2.24 g	2.24 g	--	2.24 g	--	2.24 g	--
Magnesium chloride, hexahydrate	0.45 g	0.81 g	0.81 g	--	0.81 g	--	0.81 g	--
Calcium chloride, dihydrate	0.30 g	0.52 g	0.52 g	--	0.52 g	--	0.52 g	--
Dextrose (Glucose monohydrate) (equivalent to Glucose anhydrous)	82.50 g (75.00 g)	126.5 g (115.0 g)	154.0 g (140 g)	154.0 g (140 g)	121.0 g (110.0 g)	121.0 g (110.0 g)	80.67 g (73.33 g)	80.67 g (73.33 g)

<sup>1</sup> Mixture of refined olive oil (approximately 80%) and refined soybean oil (approximately 20%) corresponding to a ratio essential fatty acids / total fatty acids of 20%.

List of excipients

Lipid emulsion chamber:

Purified egg phosphatide

Glycerol

Sodium oleate

Sodium hydroxide for pH adjustment

Nitrogen

Water for injections

Amino acid with or without electrolytes solution chamber:

Glacial acetic acid for pH adjustment

Nitrogen

Water for injections

Dextrose (glucose) with or without calcium solution chamber:

Hydrochloric acid for pH adjustment

Nitrogen

Water for injections

Packaging

The three-chamber bag is a multi-layer non-Polyvinyl Chloride bag. The inner (contact) layer of the bag material is made of a blend of polyolefinic copolymers and is compatible with amino acid solutions, dextrose solutions and lipid emulsions. Other layers are made of poly-ethylene vinyl acetate, and of a copolyester.

The dextrose chamber is fitted with an injection site to be used for addition of supplements.

The amino acids compartment is fitted with an administration site for insertion of the spike of the infusion set.

To protect from air contact, the bag is packaged in an oxygen barrier overpouch, which contains an oxygen absorber.

Pack sizes:

650 mL bag: 1 carton with 10 bags

1000 mL bag: 1 carton with 6 bags

1500 mL bag: 1 carton with 4 bags

2000 mL bag: 1 carton with 4 bags

2500 mL bag: 1 carton with 2 bags

PART II: SCIENTIFIC INFORMATION








PHARMACEUTICAL INFORMATION









Drug Substance




OLIMEL / PeriOLIMEL (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids) contains the following drug substances in the three chambers.

- Dextrose solution with or without calcium (larger outer chamber)
- Amino acid solution with or without electrolytes (sodium, potassium, magnesium, phosphate) (centre chamber)
- Lipid emulsion (a mix of refined olive oil and refined soybean oil) (smaller outer chamber)

Proper Name Chemical Name	Molecular Formula and Molecular Mass	Structural Formula	Physicochemical Properties
L-Alanine  (S)-2-aminopropionic acid	C <sub>3</sub> H <sub>7</sub> NO <sub>2</sub> 89.09		White or almost white crystalline powder or colourless crystals, freely soluble in water, very slightly soluble in alcohol.
L-Arginine  (2S)-2-amino-5-guanidinopentanoic acid	C <sub>6</sub> H <sub>14</sub> N <sub>4</sub> O <sub>2</sub> 174.20		White or almost white crystalline powder or colourless crystals, freely soluble in water, very slightly soluble in alcohol.
L-Aspartic Acid  (S)-aminosuccinic acid	C <sub>4</sub> H <sub>7</sub> NO <sub>4</sub> 133.10		White or almost white crystalline powder or colourless crystals, slightly soluble in water, practically insoluble in alcohol. It dissolves in dilute mineral acids and in dilute solutions of alkali hydroxides.
L-Glutamic Acid  (S)-2-aminoglutaric acid	C <sub>5</sub> H <sub>9</sub> NO <sub>4</sub> 147.13		White crystalline powder or colourless crystals, freely soluble in boiling water, slightly soluble in cold water, practically insoluble in acetic acid, in acetone and in alcohol.

Proper Name Chemical Name	Molecular Formula and Molecular Mass	Structural Formula	Physicochemical Properties
Glycine Aminoacetic acid	$C_2H_5NO_2$ 75.07		White or almost white crystalline powder, freely soluble in water, very slightly soluble in alcohol.
L-Histidine (S)-2-amino-1H-imidazole-4-propionic acid	$C_6H_9N_3O_2$ 155.15		White or almost white crystalline powder or colourless crystals, soluble in water, very slightly soluble in ethanol (96%).
L-Isoleucine (2S, 3S)-2-amino-3-methylpentanoic acid	$C_6H_{13}NO_2$ 131.17		White or almost white crystalline powder or flakes, sparingly soluble in water, slightly soluble in alcohol. It dissolves in dilute mineral acids and in dilute solutions of alkali hydroxides.
L-Leucine (2S)-2-amino-4-methylpentanoic acid	$C_6H_{13}NO_2$ 131.17		White or almost white crystalline powder or shiny flakes, sparingly soluble in water, practically insoluble in alcohol. It dissolves in dilute mineral acids and in dilute solutions of alkali hydroxides.
L-Lysine Acetate (2S)-2,6-diaminohexanoic acid monoacetate	$C_6H_{14}N_2O_2 \cdot C_2H_4O_2$ 206.24		White or almost white crystalline powder or colourless crystals, freely soluble in water, very slightly soluble in ethanol (96%).
L-Methionine (2S)-2-amino-4-(methylsulfanyl)butanoic acid	$C_5H_{11}NO_2S$ 149.21		White or almost white crystalline powder or colourless crystals, soluble in water, very slightly soluble in ethanol.
L-Phenylalanine (2S)-2-amino-3-phenylpropanoic acid	$C_9H_9NO_2$ 165.19		White or almost white crystalline powder or shiny, white flakes, sparingly soluble in water, very slightly soluble in alcohol. It dissolves in dilute mineral acids and in dilute solutions of alkali hydroxides.

Proper Name Chemical Name	Molecular Formula and Molecular Mass	Structural Formula	Physicochemical Properties
L-Proline  (S)-2- pyrrolidinecarboxylic acid	C <sub>5</sub> H <sub>9</sub> NO <sub>2</sub> 115.13		White or almost white crystalline powder or colourless crystals, very soluble in water, freely soluble in alcohol.
L-Serine  (S)-2-amino-3- hydroxypropionic acid	C <sub>3</sub> H <sub>7</sub> NO <sub>3</sub> 105.09		White or almost white crystalline powder or colourless crystals, freely soluble in water, practically insoluble in alcohol.
L-Threonine  (2S, 3R)-2-amino-3- hydroxybutanoic acid	C <sub>4</sub> H <sub>9</sub> NO <sub>3</sub> 119.12		White crystalline powder or colourless crystals, soluble in water, practically insoluble in ethanol.
L-Tryptophan  (2S)-2-amino-3- (indol-3-yl)propanoic acid	C <sub>11</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub> 204.23		White or almost white crystalline or amorphous powder, sparingly soluble in water, slightly soluble in alcohol. It dissolves in dilute mineral acids and in dilute solutions of alkali hydroxides.
L-Tyrosine  (S)-2-amino-3-(4- hydroxyphenyl) propionic acid	C <sub>9</sub> H <sub>11</sub> NO <sub>3</sub> 181.19		White crystalline powder or colourless crystals, very slightly soluble in water, practically insoluble in alcohol. It dissolves in dilute mineral acids and in dilute solutions of alkali hydroxides.
L-Valine  (2S)-2-amino-3- methylbutanoic acid	C <sub>5</sub> H <sub>11</sub> NO <sub>2</sub> 117.15		White or almost white crystalline powder or colourless crystals, soluble in water, very slightly soluble in ethanol.
Sodium glycerophosphate, hydrated*	C <sub>3</sub> H <sub>7</sub> Na <sub>2</sub> O <sub>6</sub> P · x H <sub>2</sub> O degree of hydration: x=4 to 6	<i>R</i> - $\alpha$ - isomer 	White crystalline powder or crystals, freely soluble in water, practically insoluble in acetone and in alcohol.
Mixture of Sodium (2 <i>RS</i> )-2,3- dihydroxypropyl phosphate and Sodium 2-hydroxy-1-	216.0 (anhydrous)	<i>S</i> - $\alpha$ - isomer 	

Proper Name Chemical Name	Molecular Formula and Molecular Mass	Structural Formula	Physicochemical Properties
(hydroxymethyl)-ethyl phosphate		$\beta$ -isomer 	
Sodium acetate trihydrate*	$C_2H_3NaO_2 \cdot 3H_2O$ 136.08		Colourless crystals, very soluble in water, soluble in alcohol.
Potassium chloride*	KCl 74.55	not provided	White or almost white crystalline powder or colourless crystals, freely soluble in water, practically insoluble in anhydrous alcohol.
Magnesium chloride hexahydrate*	$MgCl_2 \cdot 6H_2O$ 203.30	not provided	Colourless crystals, hygroscopic, very soluble in water, freely soluble in alcohol.
Calcium chloride dihydrate*	$CaCl_2 \cdot 2H_2O$ 147.01	not provided	White crystalline powder, hygroscopic, freely soluble in water, soluble in alcohol.
Dextrose D-glucose monohydrate	$C_6H_{12}O_6 \cdot H_2O$ 198.2		White crystalline powder with a sweet taste, freely soluble in water, sparingly soluble in alcohol.
Olive oil, refined	Complex mixture of triglycerides; predominant fatty acids in olive oil are oleic, palmitic and linoleic. Approximately 870 g/mol depending on the fatty acid composition.	$\begin{array}{c} CH_2-OCO-R_1 \\   \\ CH-OCO-R_2 \\   \\ CH_2-OCO-R_3 \end{array}$ <p>where <math>R_1</math>, <math>R_2</math> and <math>R_3</math> represent the fatty acids linked to the glycerol moiety of the triglyceride.</p>	Clear, colourless or greenish-yellow, transparent liquid, practically insoluble in ethanol (96%), miscible with light petroleum (50°C to 70°C). When cooled, it begins to become cloudy at 10°C and becomes a butter-like mass at about 0°C. It has a relative density of about 0.913.



Proper Name Chemical Name	Molecular Formula and Molecular Mass	Structural Formula	Physicochemical Properties
Soybean oil, refined	Complex mixture of triglycerides; predominant fatty acids in soybean oil are linoleic, oleic, palmitic and linolenic. Approximately 870 g/mol depending on the fatty acid composition.	$\begin{array}{c} \text{CH}_2\text{-OCO-R}_1 \\   \\ \text{CH-OCO-R}_2 \\   \\ \text{CH}_2\text{-OCO-R}_3 \end{array}$ where R <sub>1</sub> , R <sub>2</sub> and R <sub>3</sub> represent the fatty acids linked to the glycerol moiety of the triglyceride.	Clear, pale yellow, liquid, miscible with light petroleum (50°C to 70°C), practically insoluble in alcohol. It has a relative density of about 0.922 and a refractive index of about 1.475.

\* only contained in the formulations with electrolytes

## CLINICAL TRIALS

### Study demographics and trial design

Study ICS1063B/P01/03/Mu.F was a prospective randomized double-blind multicenter study performed to evaluate safety and nutritional efficacy of OLIMEL 5.7% (amino acids, dextrose, lipids) compared to OLICLINOMEL N8-800 (a triple chambered bag established in Europe that contains the same olive oil/soybean oil lipid, a similar amino acid profile, and dextrose). The study was conducted in a variety of patients (primarily post-surgery and trauma) who required balanced parenteral nutrition representing at least 50% of the daily nonprotein energy requirements for 5 days. The primary nutritional efficacy endpoint was transthyretin (pre-albumin) levels. Safety was evaluated using adverse events, vital signs, and biochemical markers for renal (urea, creatinine), hepatic (ASAT, ALAT, alkaline phosphatase, GGT, bilirubin), hematologic (RBC count, hemoglobin, hematocrit, platelets, WBCs, lymphocytes, neutrophils, monocytes, eosinophils, basophils) organ functions as well as glucose and lipid parameters (triglycerides, cholesterol).

Summary of patient demographics for clinical trials in specific indication					
Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
ICS1063 B/P01/03 /Mu.F	Prospective randomized double-blind multicenter controlled trial	OLIMEL 5.7%: 29 kcal/kg/day; protein 1.5 g/kg/day; 5 days  OLICLINOMEL: 29 kcal/kg/day; protein 1.5 g/kg/day ; 5 days  via central vein	ITT population OLIMEL 5.7%: n=24 OLICLINOMEL : n=26	OLIMEL 5.7%: 56±15 years OLICLINOMEL: 52±21 years	OLIMEL 5.7%: 17 male, 7 female OLICLINOMEL: 16 male, 10 female

## Study results

### Results of study ICS1063B/P01/03/Mu.F in specific indication

Primary Endpoints	Associated value and statistical significance for Drug at specific dosages	Associated value and statistical significance for Placebo or active control
Transthyretin levels	No differences were found between the two nutritional groups for transthyretin levels. Day 1 to Day 5: 144±75 mg/L to 206±142 mg/L	Day 1 to day 5: 146±83 mg/L to 181±82 mg/L

Adverse events were similar between groups (see [Clinical Trial Adverse Drug Reactions](#)). There were no differences between groups for glucose, lipids, renal function, hepatic function, or hematologic function.

## DETAILED PHARMACOLOGY

No studies were performed.

## TOXICOLOGY

Studies on the carcinogenic potential, reproductive and developmental toxicity, and genotoxic potential have not been performed on the finished product for OLIMEL/ PeriOLIMEL (amino acids WITH electrolytes, dextrose, lipids) or (amino acids, dextrose, lipids).

## Amino Acid Compartment

The amino acid solution with electrolytes contains sodium glycerophosphate as a source of nutritional phosphorus. Single-dose and 28-day repeated dose toxicity studies have been performed in rats and dogs with sodium glycerophosphate.

### **Single-dose Toxicity**

Rats received a single intravenous bolus infusion of sodium glycerophosphate at doses of 625, 1,250 or 2,500 mg/kg. At the 2500 mg/kg dose, 10% mortality and clinical signs of toxicity were observed. Therefore, 1250 mg/kg was identified as the maximum tolerated dose following intravenous infusion in rats.

Dogs were given a single intravenous bolus infusion at doses of 625, 1250 or 1875 mg/kg. At the 1875 mg/kg dose, vomiting, jaw discomfort, half-closed eyes, loss of balance and sneezing were observed during, or just after dosing. Vomiting was noted during infusion in the 1250 mg/kg group. The no observed effect level was considered to be 625 mg/kg/day.

### **Repeat-Dose Toxicity**

Rats were given 312.5, 625, or 1250 mg/kg of sodium glycerophosphate intravenously daily for 28 consecutive days. No test article-related changes were observed.

Dogs received sodium glycerophosphate by daily intravenous infusion at the dose-levels of 312.5, 625 or 1250 mg/kg/day for 28 consecutive days. At 1250 mg/kg/day, vomiting, rubbing of the head on the floor, shaking head, scratching of ear(s), half-closed eyes, loss of balance, sitting position and lateral recumbency were observed primarily during the first two days of the treatment period. A slight prolongation of the QT and QTc interval durations was noted in females given 1250 mg/kg/day throughout the treatment period. An increased QT interval at the 1250 mg/kg dose could be attributed to a transient, localized hypocalcemia associated with the rapid administration rate (compared to the therapeutic rate of sodium glycerophosphate delivery in humans). Sodium glycerophosphate was clinically well tolerated and did not cause any laboratory or histopathological changes at the indicated doses.

### **Lipid Compartment**

#### **Single-dose Toxicity**

Single-dose toxicity was investigated in the mouse and rat to compare the LD<sub>50</sub> of the lipid emulsion in OLIMEL / PeriOLIMEL with that of 20% soybean based emulsions.

LD<sub>50</sub> values were comparable at around 100-112 mL/kg (corresponding to approximately 20 g lipid/kg) in both species with rapid infusion.

#### **Repeat-Dose Toxicity**

The lipid emulsion in OLIMEL / PeriOLIMEL was administered to rats and dogs by intravenous infusion in studies lasting up to 91 days. The key studies conducted and the notable findings are presented in the table below.

**Repeated Dose Toxicity Studies**

<b>Type of Study</b>	<b>Species and Strain</b>	<b>Method of Administration</b>	<b>Duration of Dosing</b>	<b>Doses (mL/kg/day)</b>	<b>Key Findings</b>
30 Day toxicity study in the rat	Rat Sprague Dawley CD (Charles River)	IV infusion	30 days	90 at a rate of 1.2 mL/kg/min	Hematuria Decreased food consumption Regenerative anemia Mild thrombocytopenia Increased hepatic enzymes Increased serum cholesterol and phospholipids Hepatocellular vacuolation and necrosis Hepatic and splenic macrophage pigmentation Splenic macrophage vacuolation Interstitial and tubular nephritis
30 Day toxicity study in the rat	Rat Sprague Dawley CD (Charles River)	IV infusion	30 days	75 at a rate of 1.5 mL/kg/min	Hematuria Decreased food consumption Regenerative anemia Mild thrombocytopenia Increased hepatic enzymes Increased serum phospholipids Hepatic vacuolation and inflammation Hepatic and splenic macrophage pigmentation Splenic macrophage vacuolation Interstitial and tubular nephritis
90 Day toxicity study in the rat	Rat Sprague Dawley CD (Charles River)	IV infusion	90 days	15, 30 and 60 at a rate of 2 mL/kg/min	Hematuria Decreased food consumption Mild anemia at 60 mL/kg Dose-dependent hepatic vacuolation and inflammation Hepatic and splenic macrophage pigmentation Splenic macrophage vacuolation

<b>Type of Study</b>	<b>Species and Strain</b>	<b>Method of Administration</b>	<b>Duration of Dosing</b>	<b>Doses (mL/kg/day)</b>	<b>Key Findings</b>
30 Day toxicity study in the dog	Dog Beagle	IV infusion	30 days	45 at a rate of 0.2 mL/kg/min	Decreased food consumption Increased hepatic enzymes Increased serum cholesterol and phospholipids Hepatocellular vacuolation Hepatic and splenic macrophage pigmentation Splenic macrophage vacuolation
30 Day toxicity study in the dog	Dog Beagle	IV infusion	30 days	60 at a rate of 0.2 mL/kg/min	Decreased food consumption Regenerative anemia Mild thrombocytopenia Increased hepatic enzymes Increased serum phospholipids Hepatic vacuolation and inflammation, Hepatic and splenic macrophage pigmentation Splenic macrophage vacuolation Renal tubular vacuolation
30 Day toxicity study in the dog	Dog Beagle	IV infusion	30 days	60 at a rate of 0.2 mL/kg/min	Decreased food consumption Regenerative anemia Mild thrombocytopenia Increased hepatic enzymes Increased serum phospholipids Hepatic vacuolation and inflammation, Hepatic and splenic macrophage pigmentation Splenic macrophage vacuolation Renal tubular vacuolation
91 Day toxicity study in the dog	Dog Beagle	IV infusion	91 days	15, 22.5, 30 0.2 mL/kg/min	Decreased food consumption Mild anemia at 30 mL/kg Increased hepatic enzymes at $\geq 22.5$ mL/kg Increased serum cholesterol and phospholipids Hepatocellular vacuolation at 30 mL/kg Hepatic and splenic macrophage pigmentation Splenic macrophage vacuolation

## REFERENCES

1. Allwood MC. Reflections on current issues concerning the stability of parenteral nutrition mixtures. *Nutrition* 2002, Jul-Aug; 18(7-8):691-2.
2. Anker SD, Laviano A, Filippatos G, John M, Paccagnella A, Ponikowski P, et al. ESPEN guidelines on parenteral nutrition: on cardiology and pneumology. *Clin Nutr.* 2009 28:455–60.
3. ASPEN. Guidelines for the USE of Parenteral and Enteral Nutrition in Adult and Pediatric Patients. *JPEN* 2002; 26 (supplement): 1-138.
4. The ASPEN Nutrition Support Core Curriculum. American Society of Parenteral and Enteral Nutrition. 2007, 1-824.
5. Baxter Corporation. CLINOLEIC 20% Product Monograph (lipid emulsion in OLIMEL).
6. Berger M, Chioloro R. Chapter 15: Trauma and burns. In: Rombeau JL, Caldwell MD, editors. *Clinical nutrition – parenteral nutrition*. Philadelphia: WB Saunders Company; 1993.
7. Booth SL, Suttie JW. Dietary intake and adequacy of vitamin K. *J Nutr.* 1998 May;128(5):785-8.
8. Bozzetti F, Arends J, Lundholm K, Micklewright A, Zurcher Z, Muscaritoli M, et al. ESPEN guidelines on parenteral nutrition: non-surgical oncology. *Clin Nutr.* 2009a 28:445–54.
9. Bozzetti F, Forbes A. ESPEN guidelines on parenteral nutrition: present status and perspectives for future research. *Clin Nutr.* 2009b 28:359–64.
10. Braga M, Ljungqvist O, Soeters P, Fearon K, Weimann A, Bozzetti F. ESPEN Guidelines on Parenteral Nutrition: surgery ESPEN guidelines on parenteral nutrition: surgery. *Clin Nutr.* 2009 28: 378-86.
11. Cane RD, Harrison RA, Shapiro BA, Kavanaugh J. The spectrophotometric absorbance of Intralipid. *Anesthesiology* 1980, 53(1):53-55
12. Cano et al. ESPEN Guidelines on Parenteral Nutrition: Adult Renal Failure. *Clin Nutr* 2009; 28(4):401-414.
13. Carlin A, Walker WA. Rapid development of vitamin K deficiency in an adolescent boy receiving total parenteral nutrition following bone marrow transplantation. *Nutr Rev.* 1991 Jun;49(6):179-83.

14. Chakraverty R, Davidson S, Peggs K, Stross P, Garrard C, Littlewood TJ. The incidence and cause of coagulopathies in an intensive care population. *Br J Haematol*. 1996 May;93(2):460-3.
15. Denet A-R. (2006). Supplementation of OliClinisol N4-600E 1L and N9-XXXE 1L, Baxter Study 787-RF-NIV, 1–14.
16. Duerksen DR, Papineau N. The prevalence of coagulation abnormalities in hospitalized patients receiving lipid-based parenteral nutrition. *JPEN*. 2004 Jan-Feb;28(1):30-33.
17. Gianotti et al., ESPEN guidelines on parenteral nutrition: pancreas. *Clin Nutr* 2009; 28:428-435.
18. Grafmeyer D, Bondon M, Manchon M, Levillain P. The influence of bilirubin, haemolysis and turbidity on 20 analytical tests performed on automatic analysers. Results of an interlaboratory study. *Eur J Clin Chem Clin Biochem* 1995, 33(1):31-52.
19. Hill SE, Heldman LS, Goo ED, Whippo PE, Perkinson JC. Fatal microvascular pulmonary emboli from precipitation of total nutrient admixture solution. *JPEN J Parenter Enteral Nutr* 1996 Jan-Feb;20(1):81-87
20. Howdieshell TR, Sussman A, Dipiro J, McCarten M, Mansberger A Jr. Reliability of in vivo mixed venous oximetry during experimental hypertriglyceridemia. *Crit Care Med* 1992, 20(7):999-1004.
21. Koletzko B, Goulet O, Hunt J, et al. Guidelines on Paediatric Parenteral Nutrition of the European Society of Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) and the European Society for Clinical Nutrition and Metabolism (ESPEN), Supported by the European Society of Paediatric Research (ESPR). *J Pediatr Gastroenterol Nutr* 2005;41(Suppl 2):S1–87.
22. Martinez MJ, Folch N, Torregrosa N, Alfaro A, Obaldia MC. Coagulation and total parenteral nutrition: the effect of 2 lipid emulsions. (Article in Spanish.) *Nutr Hosp*. 1995 Mar-Apr;10(2):74-80.
23. McKinnon BT. FDA safety alert: hazards of precipitation associated with parenteral nutrition. *Nutr Clin Pract* 1996 Apr, 11(2):59-65.
24. McNearney T, Bajaj C, Boyars M, Cottingham J, Haque A. Total parenteral nutrition associated crystalline precipitates resulting in pulmonary artery occlusions and alveolar granulomas. *Dig Dis Sci* 2003 Jul;48(7):1352-1354
25. Mirtallo J, Canada T, Johnson D, Kumpf V, Petersen C, Sacks G, Seres D, Guenter P. Task Force for the Revision of Safe Practices for Parenteral

- Nutrition, Special Report: Safe practices for parenteral nutrition. *JPEN J Parenter Enteral Nutr.* 2004 Nov-Dec;28 (6 Suppl):S39-70.
26. Miyada D, Tipper P, Jantsch D, Simpkins H, Steele W, Flores O. The effect of hyperlipidemia on Technicon SMAC measurements. *Clin Biochem* 1982, 15(4):185-8.
  27. Payne-James J, Grimble GK, Silk DBA: Artificial nutrition Support in Clinical Practice, 2<sup>nd</sup> ed, Greenwich Medical Media Limited, London, 2001, 1-798.
  28. Pittiruti M, Hamilton H, Biffi R, MacFie J, Pertkiewicz Merk. ESPEN guidelines on parenteral nutrition: central venous catheters (access, care, diagnosis and therapy of complications). *Clin Nutr.* 2009 28(1):365–77.
  29. Plauth M, Cabré E, Campillo B, Kondrup J, Marchesini G, Schütz T, et al. ESPEN Guidelines on Parenteral Nutrition: hepatology. *Clin Nutr.* 2009 Aug;28(4):436-44.
  30. Powell-Tuck J, Gosling P, Lobo DN, Allison SP, Carlson GL, Gore M, et al. British consensus guidelines on intravenous fluid therapy for adult surgical patients. *NHS Evidence.* Oct 2008. Obtained from website on 2 Feb 2010: [http://www.ics.ac.uk/intensive\\_care\\_professional/british\\_consensus\\_guidelines\\_on\\_intravenous\\_fluid\\_therapy\\_for\\_adult\\_surgical\\_patients\\_\\_giftasup\\_\\_2008](http://www.ics.ac.uk/intensive_care_professional/british_consensus_guidelines_on_intravenous_fluid_therapy_for_adult_surgical_patients__giftasup__2008)
  31. Pugh MB et al., ed. 2006. Stedman's Medical Dictionary. Baltimore, MD. Lippincott Williams & Wilkins. ISBN 1071-3964. STAT!Ref Online.
  32. Randall AG, Garcia-Webb P, Beilby JP. Interference by haemolysis, icterus and lipaemia in assays on the Beckman Synchron CX5 and methods for correction. *Ann Clin Biochem* 1990, 27(Pt 4):345-52.
  33. Reedy JS, Kuhlman JE, Voytovich M. Microvascular pulmonary emboli secondary to precipitated crystals in a patient receiving total parenteral nutrition: a case report and description of the high-resolution CT findings. *Chest* 1999 Mar;115(3):892-895
  34. Robinson LA, Wright BT. Central venous catheter occlusion caused by body-heat-mediated calcium phosphate precipitation. *Am J Hosp Pharm* 1982 Jan, 39(1):120-121
  35. Robinson RG, Fawcett CA, van Ru AM. Multiple central venous line occlusions following an error in parenteral nutrition formulation. *Aust NZ J Surg* 1986;56:347-349



36. Sehgal LR, Sehgal HL, Rosen AL, Gould SA, Moss GS. Effect of Intralipid on measurements of total hemoglobin and oxyhemoglobin in whole blood. *Crit Care Med* 1984, 12(10):907-9.
37. Shay DK, Fann LM, Jarvis WR. Respiratory distress and sudden death associated with receipt of a peripheral parenteral nutrition admixture. *Infect Control Hosp Epidemiol* 1997;18(12):814-817
38. Singer P, Berger M, Van den Berghe G, Biolo G, Calder P, Forbes A, et al. ESPEN guidelines on parenteral nutrition: intensive care. *Clin Nutr.* 2009 28:387–400
39. Staun M, Pironi L, Bozzetti F, Baxter J, Forbes A, Joly F, et al. ESPEN Guidelines on parenteral nutrition: home parenteral nutrition. *Clin Nutr.* 2009 28:467–79.
40. Stennett DJ, Gerwick WH, Egging PK, Christensen JM. Precipitate analysis from an indwelling total parenteral nutrition catheter. *JPEN J Parenter Enteral Nutr* 1988, 12(1):88-92.
41. Sutton BJ, Gould HJ. The human IgE network. *Nature.* 1993 Dec 2;366(6454):421-8.
42. U.S. Dept. of Health and Human Services, U.S. Food and Drug Administration. FDA Safety Alert: Hazards of precipitation associated with parenteral nutrition. Issued April 18, 1994. Available at: <http://www.fda.gov/MedicalDevices/Safety/AlertsandNotices/PublicHealthNotifications/ucm238205.htm>
43. Van Gossum A, Cabre E, Hebuterne X, Jeppesen P, Krznaric Z, Messing B, et al. ESPEN Guidelines on parenteral nutrition: gastroenterology. *Clin Nutr.* 2009 28:415-27.

## IMPORTANT: PLEASE READ

### READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

#### PART III: CONSUMER INFORMATION

##### **OLIMEL 3.3% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
3.3% w/v & 0.7% w/v, 11.5% w/v, 4% w/v

##### **OLIMEL 4.4%**

Amino acids, dextrose, lipids  
Injectable Emulsion  
4.4% w/v, 14% w/v, 4% w/v

##### **OLIMEL 4.4% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
4.4% w/v & 0.7% w/v, 14% w/v, 4% w/v

##### **OLIMEL 5.7%**

Amino acids, dextrose, lipids  
Injectable Emulsion  
5.7% w/v, 11% w/v, 4% w/v

##### **OLIMEL 5.7% E**

Electrolytes  
Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
5.7% w/v & 0.7% w/v, 11% w/v, 4% w/v

##### **OLIMEL 7.6%**

Amino acids, dextrose, lipids  
Injectable Emulsion  
7.6% w/v, 7.3% w/v, 3.5% w/v

##### **OLIMEL 7.6% E**

Electrolytes  
Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
7.6% w/v & 0.8% w/v, 7.3% w/v, 3.5% w/v

##### **PeriOLIMEL 2.5% E**

Amino acids **WITH** electrolytes, dextrose, lipids  
Injectable Emulsion  
2.5% w/v & 0.4% w/v, 7.5% w/v, 3% w/v

**Read this carefully before you start taking OLIMEL / PeriOLIMEL product and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about OLIMEL / PeriOLIMEL product.**

#### **ABOUT THIS MEDICATION**

##### **What the medication is used for:**

OLIMEL / PeriOLIMEL product is used to provide nutrition to adults through a tube into a vein when normal feeding by mouth is not possible or suitable.

OLIMEL / PeriOLIMEL product must only be used under medical supervision.

##### **What it does:**

The use of OLIMEL / PeriOLIMEL product is a way to ensure that patients who are unable to eat get an adequate intake of energy, nitrogen and other nutrients, and helps to treat or prevent malnutrition.

##### **When it should not be used:**

Do not use OLIMEL / PeriOLIMEL product if:

- You are allergic to any ingredients (such as egg, soybean proteins, olive products.) (See **What the medicinal ingredients are** and **What the important nonmedicinal ingredients are**.)
- You have liver failure or coma resulting from liver failure.
- You have kidney failure and are not on dialysis
- You have an allergy to corn or corn products since this product contains dextrose from corn.
- Your body has problems processing certain amino acids that are included in OLIMEL / PeriOLIMEL product.
- Your body has severe problems metabolizing (breaking down) fat.
- You have especially high levels of fats in your blood.
- You have acute pancreatitis (severe inflammation of pancreas) in association with hyperlipidemia (high blood fat levels).
- You have hyperglycemia (too much sugar in your blood), which is not controlled.
- You have high plasma concentrations of one of the electrolytes included in some of the OLIMEL and PeriOLIMEL product.
- You have a disorder resulting in high blood levels of substances such as potassium (hyperkalemia), calcium (hypercalcemia), phosphorus (hyperphosphatemia), sodium (hypernatremia) and magnesium (hypermagnesemia).
- You are being simultaneously administered ceftriaxone with intravenous calcium-containing solutions, including OLIMEL / PeriOLIMEL product, through the same infusion line.

##### **What the medicinal ingredients are:**

OLIMEL / PeriOLIMEL product is an emulsion for infusion, supplied in a bag with three chambers.

one chamber contains a solution of 17 amino acids (L-Alanine, L-Arginine, L-Aspartic Acid, L-Glutamic Acid, Glycine, L-Histidine, L-Isoleucine, L-Leucine, L-Lysine Acetate, L-Methionine, L-Phenylalanine, L-Proline, Serine, L-Threonine, L-Tryptophan, L-Tyrosine, L-Valine) with or without electrolytes (sodium, potassium, magnesium, phosphate); one chamber contains a dextrose solution with or without calcium; one chamber contains a lipid emulsion, which is a mix of refined olive oil and refined soybean oil.

**What the nonmedicinal ingredients are:**

Glacial acetic acid (for pH adjustment)  
Glycerol  
Hydrochloric acid (for pH adjustment)  
Nitrogen  
Purified egg phosphatide  
Sodium hydroxide (for pH adjustment)  
Sodium oleate  
Water for injection

**What dosage forms it comes in:**

OLIMEL 3.3% E (3.3% w/v Amino acids with 0.7% w/v electrolytes, 11.5 % dextrose, 4% w/v lipids) Injectable Emulsion  
OLIMEL 4.4% (4.4% w/v Amino acids, 14% w/v dextrose, 4% w/v lipids) Injectable Emulsion  
OLIMEL 4.4% E (4.4% w/v Amino acids with 0.7% w/v electrolytes, 14% w/v dextrose, 4% w/v lipids) Injectable Emulsion  
OLIMEL 5.7% (5.7% w/v Amino acids, 11% w/v dextrose, 4% w/v lipids) Injectable Emulsion  
OLIMEL 5.7% E (5.7% w/v Amino acids with 0.7% w/v electrolytes, 11% w/v dextrose, 4% w/v lipids) Injectable Emulsion  
OLIMEL 7.6 % (7.6% w/v Amino acids, 7.3% w/v dextrose, 3.5% w/v lipids) Injectable Emulsion  
OLIMEL 7.6% E (7.6% w/v Amino acids with 0.8% w/v electrolytes, 7.3% w/v dextrose, 3.5% w/v lipids) Injectable Emulsion  
PeriOLIMEL 2.5% E (2.5% w/v Amino acids with 0.4% w/v electrolytes, 7.5% w/v dextrose, 3% w/v lipids) Injectable Emulsion

**WARNINGS AND PRECAUTIONS**

**BEFORE you use OLIMEL / PeriOLIMEL talk to your doctor or pharmacist if you:**

- are allergic to any ingredients (such as egg, soybean proteins, olive products.) (See **What the medicinal ingredients are** and **What the nonmedicinal ingredients are**.)
- suffer from metabolic acidosis (when the blood is excessively acid)
- have kidney or liver problems

- have acute respiratory distress syndrome
- are taking any other medicines on a regular basis
- are pregnant or intend to become pregnant
- are breastfeeding or intend to breastfeed
- have hereditary fructose intolerance as this product may contain small amounts of fructose

Sometimes treatment with this medicine is not suitable, such as if you have:

- Acute pulmonary oedema (collection of fluid into the lung tissue), hyperhydration (too much water in your body), heart failure, dehydration, untreated diabetes mellitus, severe metabolic acidosis (when the blood is excessively acid), severe disorders of fat metabolism, liver disease, or kidney disease. You may not be able to tell if you have some of these conditions, so be sure to check with your doctor first.
- Precipitates (e.g. solid particles) can appear in the OLIMEL / PeriOLIMEL formulation when a calcium-containing product is added to OLIMEL / PeriOLIMEL formulation. It has been reported that these injected precipitates can block an artery in the lungs, cause breathing difficulties and even death. Your doctor will decide with caution, whether a calcium-containing product will be mixed with OLIMEL / PeriOLIMEL product.

In addition to inspection of the solution or emulsion, the infusion set and catheter should also periodically be checked for precipitates.

If you have difficulties breathing, stop the infusion and contact your healthcare professional immediately.

In all cases, your doctor will base his/her decision to give you this medicine on factors such as age, weight and clinical condition, together with the results of any tests that he/she has performed. Always be sure to check with your doctor if anything about your condition changes.

Your doctor will need to monitor how you are doing while you are on this medicine. This means that you will need to have laboratory tests done on a routine basis.

**INTERACTIONS WITH THIS MEDICATION**

No interaction studies have been performed.

Soybean oil in OLIMEL / PeriOLIMEL product has a natural content of vitamin K that may counteract the blood-thinning (anticoagulant) activity of anticoagulants such as

warfarin. Discuss with your doctor if you are taking warfarin.

There may also be an interaction between the OLIMEL / PeriOLIMEL products containing electrolytes and the antibiotic ceftriaxone.

OLIMEL / PeriOLIMEL product must NOT be administered simultaneously with blood through the same infusion tubing.

There may be interactions between the nutrients in OLIMEL / PeriOLIMEL product and one or more of your medications, for example blood thinners. You should review your medications with your doctor.

### PROPER USE OF THIS MEDICATION

OLIMEL / PeriOLIMEL product can be given in a hospital or managed care facility, or at home under the supervision of a doctor or other health care professional.

After appropriate training and with the agreement of your medical team, you may be able to administer the product yourself. Your doctor's instructions must be followed exactly when taking OLIMEL / PeriOLIMEL product.

Before using the product, the bag must be prepared as shown in the pictures.

Use only if the solutions are clear, colourless or slightly yellow, practically free of visible particles and if the emulsion is homogeneous and milk-like.

Use only if the bag is not damaged and if the non-permanent seals between the chambers are intact (no mixing of the 3 chambers contents).


Make sure the product is at room temperature.

Aseptic conditions must be followed (cleaning of hands).

OLIMEL product must be delivered through a central vein. PeriOLIMEL 2.5% E may be delivered through a central or peripheral vein.

Your healthcare professional will provide instructions on the preparation of the site and route of administration (central or peripheral vein).

OLIMEL / PeriOLIMEL product should only be used once. Discard unused portion, do not use a partially used bag.



### Usual adult dose:

Your doctor will select PeriOLIMEL product or the best OLIMEL product for you, based on your body weight. He/she will also consider how much energy you need and how well your body can handle the amino acids, lipids, and electrolytes in the different solutions

Your doctor will also specify a flow rate corresponding to your needs and medical condition.

Always use OLIMEL / PeriOLIMEL product exactly as your doctor has told you to. You should check with your doctor if you are not sure.

The product infusion may continue for as long as the doctor advises, depending upon your medical condition. The infusion of one triple-chamber bag usually lasts between 12 and 24 hours. The maximum daily dose should not be exceeded.

### Overdose:

If your dose is too high or is infused too quickly, the amino acid content may make your blood too acid, the dextrose content may increase the glucose in your blood and urine, or the lipid content may increase the fats in your blood. Giving too high a volume may cause nausea, vomiting, headache, hot flush, excessive sweating and shivering.

In some severe cases, your doctor may have to give you temporary renal dialysis to help your kidneys eliminate any excess nutrients.

To prevent these events occurring, your doctor will regularly monitor your condition and test your blood and urine parameters.

If you feel you have been given too much or have taken too much OLIMEL / PeriOLIMEL product, contact your healthcare practitioner (e.g. doctor), hospital emergency department or the regional poison control centre, even if there are no symptoms.

**Missed Dose:**

If you miss or forget to take one or more doses of OLIMEL / PeriOLIMEL product, contact your doctor as soon as possible. Your doctor will instruct you about how to re-start your treatment and what flow rate to use.

DO NOT take a double dose to make up for forgotten individual doses.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

If you notice any changes in the way you feel during or after the treatment, tell your doctor or another member of your medical team immediately.

OLIMEL / PeriOLIMEL can cause abnormal blood test results. Your doctor will decide when to perform blood tests and will interpret the results.

If any side effect gets serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or a member of your medical team right away.

Side effects may include nausea, diarrhea and abdominal pain.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and seek immediate medical help
		Only if severe	In all cases	
<b>Uncommon</b>	<b>Allergic Reaction:</b> fever,			√

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and seek immediate medical help
		Only if severe	In all cases	
	shivering, skin rashes, breathing difficulties, severe headache			
<b>Rare</b>	<b>Liver Problems:</b> increased levels of liver enzymes, yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite, increase in size of the liver and spleen		√	
	<b>Decreased Platelets:</b> bruising, bleeding, fatigue and weakness		√	
	<b>Fat Overload Syndrome:</b> too much fat in the blood, fever, jaundice (yellowing of the skin and eyes), anemia (low red blood cells), low white blood cells and blood platelets, increase in size of the spleen and liver problems with blood clotting,			√

## SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and seek immediate medical help
		Only if severe	In all cases	
	trouble breathing, coma			
	<b>Precipitates in the Blood Vessels Leading to or from the Lungs:</b> difficulty breathing, shortness of breath, chest pain, coughing			√
	<b>Injection Site Reaction:</b> pain, redness and/or swelling at the injection site		√	

*This is not a complete list of side effects. For any unexpected effects while taking OLIMEL / PeriOLIMEL product, contact your doctor or a member of your medical team or pharmacist.*

### HOW TO STORE IT

Do not freeze. Store the unmixed product in the overpouch at 15°C to 30°C.

Do not use OLIMEL / PeriOLIMEL product after the expiry date which is printed on the container and the outer packaging (MM/YYYY). The expiry date refers to the last day of that month.

Once the seals between the three chambers have been broken and the product has been mixed, the product should be used **immediately**.

**This medicine must be at room temperature to be administered.**

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Keep out of reach and sight of children.

### REPORTING SIDE EFFECTS

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345

*NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance program does not provide medical advice.*

### MORE INFORMATION

If you want more information about OLIMEL/ PeriOLIMEL:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada.html>); the Baxter website (Baxter.ca), or by calling 1-888-719-9955.

This leaflet was prepared by Baxter Corporation, Mississauga, Ontario L5N 0C2, Canada

Baxter, Olimel, and Periolumel are trademarks of Baxter International Inc.

Last revised: May 20, 2020